



This is a repository copy of *The Solar Activity Monitor Network - SAMNet*.

White Rose Research Online URL for this paper:
<https://eprints.whiterose.ac.uk/175746/>

Version: Accepted Version

Article:

Erdelyi, R. orcid.org/0000-0003-3439-4127, Brigitta Korsos, M., Huang, X. et al. (37 more authors) (2021) The Solar Activity Monitor Network - SAMNet. *Journal of Space Weather and Space Climate*. ISSN 2115-7251

<https://doi.org/10.1051/swsc/2021025>

© 2021 The author(s). This is an author-produced version of a paper subsequently published in *Journal of Space Weather and Space Climate*. Uploaded in accordance with the publisher's self-archiving policy.

Reuse

Items deposited in White Rose Research Online are protected by copyright, with all rights reserved unless indicated otherwise. They may be downloaded and/or printed for private study, or other acts as permitted by national copyright laws. The publisher or other rights holders may allow further reproduction and re-use of the full text version. This is indicated by the licence information on the White Rose Research Online record for the item.

Takedown

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing eprints@whiterose.ac.uk including the URL of the record and the reason for the withdrawal request.



eprints@whiterose.ac.uk
<https://eprints.whiterose.ac.uk/>

The Solar Activity Monitor Network - SAMNet

Robertus Erdélyi^{1,2,3}, Marianna B. Korsós^{4,2,3}, Xin Huang⁵, Yong Yang⁶, Danielle Pizzey⁷, Steven A. Wrathmall⁷, Ifan G. Hughes⁷, Martin J. Dyer⁸, Vikram S. Dhillon^{8,9}, Bernadett Belucz^{2,3}, Roman Brajša¹⁰, Piyali Chatterjee¹¹, Xuewu Cheng⁶, Yuanyong Deng⁵, Santiago Vargas Domínguez¹², Raúl Joya¹³, Peter Gömöry¹⁴, Norbert G. Gyenge^{1,3}, Arnold Hanslmeier¹⁵, Ales Kucera¹⁴, David Kuridze^{4,16}, Faquan Li⁶, Zhong Liu^{17,18}, Long Xu⁵, Mihalis Mathioudakis¹⁹, Sarah Matthews²⁰, R. T. James McAteer^{21,22}, Alexei A. Pevtsov^{23,24}, Werner Pötzi²⁵, Paolo Romano²⁶, Jinhua Shen²⁷, János Temesváry³, Andrey G. Tlatov²⁸, Charles Triana²⁹, Dominik Utz^{15,30}, Astrid M. Veronig^{15,25}, Yuming Wang^{31,32}, Yihua Yan^{5,33}, Teimuraz Zaqarashvili^{15,16,34}, Francesca Zuccarello³⁵

1. Solar Physics & Space Plasma Research Center (SP2RC), School of Mathematics and Statistics, University of Sheffield, Hicks Building, Hounsfield Road, S3 7RH, UK
2. Dep. of Astronomy, Eötvös Loránd Univ., Pázmány P. sétány 1/A, Budapest, H-1117, Hungary
3. Gyula Bay Zoltán Solar Observatory (GSO), Hungarian Solar Physics Foundation (HSPF), Petőfi tér 3., Gyula, H-5700, Hungary
4. Department of Physics, Aberystwyth University, Ceredigion, Cymru, SY23 3BZ, UK
5. CAS Key Laboratory of Solar Activity, National Astronomical Observatories, Chinese Academy of Sciences, A20 Datun Rd. Chaoyang District, Beijing 100012, China
6. State Key Laboratory of Magnetic Resonance and Atomic and Molecular Physics, Innovation Academy for Precision Measurement Science and Technology, Chinese Academy of Science (CAS), West No.30 Xiao Hong Shan, Wuhan 430071 China
7. Joint Quantum Centre (JQC) Durham-Newcastle, Department of Physics, Durham University, South Road, Durham, DH1 3LE, UK
8. Department of Physics and Astronomy, University of Sheffield, Sheffield, S3 7RH, UK
9. Instituto de Astrofísica de Canarias, E-38205 La Laguna, Tenerife, Spain
10. Hvar Observatory, Faculty of Geodesy, Univ. of Zagreb, Kačićeva 26, 10000 Zagreb, Croatia
11. Indian Institute of Astrophysics (IIA), 2nd Block Koramangala Bangalore-560034, India
12. Observatorio Astronómico Nacional, Universidad Nacional de Colombia, Bogotá, Colombia
13. Observatorio Astronómico, Universidad Sergio Arboleda, Bogotá, Colombia
14. Astronomical Institute, Slovak Academy of Sciences, 05960 Tatranská Lomnica, Slovakia
15. Institute of Physics, University of Graz, Universitätsplatz 5, A-8010 Graz, Austria
16. Abastumani Astrophysical Observatory, Mount Kanobili, Georgia
17. Yunnan Observatories, Chinese Academy of Sciences, Kunming 650011, China
18. School of Astronomy and Space Science, Univ. of CAS, Beijing 100049, China
19. Astrophysics Research Centre, School of Mathematics and Physics, Queen's University Belfast, Belfast BT7 1NN, UK
20. UCL-Mullard Space Science Lab., Holmbury St Mary, Dorking, Surrey, RH5 6NT, UK

21. Dep. of Astronomy, New Mexico State Univ, P.O. Box 30001, MSC 4500, Las Cruces, NM 88003, USA
22. Sunspot Solar Observatory, Sunspot, NM 88349, USA
23. National Solar Observatory, Boulder, CO 80303, USA
24. Pulkovo Astronomical Observatory, Russian Academy of Sciences, Saint Petersburg, 196140, Russian Federation
25. Kanzelhöhe Observatory for Solar and Environmental Research, University of Graz, Kanzelhöhe 19, A-9521 Treffen, Austria
26. INAF Osservatorio Astrofisico di Catania, Via S. Sofia 78, I 95123 Catania, Italy
27. Xinjiang Astromonomical Observatory, CAS, 150 Science 1-Street, Urumqi, 830011, China
28. Kislovodsk Solar Mountain Astronomical Station, Central Astronomical Observatory of the Russian Academy of Sciences, Kislovodsk, 357700 Russia
29. Observatorio AstroExplor, Boyacá, Colombia
30. Instituto de Astrofísica de Andalucía, Granada, Spain
31. CAS Key Laboratory of Geospace Environment, School of Earth and Space Sciences, University of Science and Technology of China, Hefei 230026, China
32. CAS Center for Excellence in Comparative Planetology, USTC, Hefei 230026, China
33. School of Astronomy and Space Science, Univ. of Chinese Acad. of Sciences, Beijing, China
34. School of Natural Sciences and Medicine, Ilia State Univ., Cholokashvili ave. 3/5, Tbilisi, Georgia
35. Dipartimento di Fisica e Astronomia “Ettore Majorana”, Università di Catania, Via S. Sofia 78, I-95123 Catania, Italy

ABSTRACT

The Solar Activity Magnetic Monitor (SAMM) Network (SAMNet) is a future UK-led international network of ground-based solar telescope stations. SAMNet, at its full capacity, will continuously monitor the Sun's intensity, magnetic and Doppler velocity fields at multiple heights in the solar atmosphere (from photosphere to upper chromosphere). Each SAMM sentinel will be equipped with a cluster of identical telescopes each with different magneto-optical filter (MOFs) to take observations in K I, Na D and Ca I spectral bands. A subset of SAMM stations will have white-light coronagraphs and emission line coronal spectropolarimeters. The objectives of SAMNet are to provide observational data for the space weather research and forecast. The goal is to achieve an operationally sufficient lead time of e.g. flare warning of 2-8 hours, and provide much sought-after continuous synoptic maps (e.g., LoS magnetic and velocity fields, intensity) of the lower solar atmosphere with a spatial resolution limited only by seeing or diffraction limit, and with a cadence of 10 minutes. The individual SAMM sentinels will be connected into their master HQ hub where data received from all the slave stations will be automatically processed and flare warning issued up to 26 hrs in advance.

Key words. Sun-flares-precursors

1. Introduction

The Earth's atmosphere and the magnetosphere are constantly buffeted by a continuous flow of solar particles originating in the solar atmosphere. This continuous flow may suddenly be enhanced and

enter into the Earth's upper atmosphere when a high-intensity solar flare (e.g. above M5-class, using the GOES classification) or a massive coronal mass ejection (CME) erupts in the solar atmosphere and reaches Earth after propagating across the interplanetary space. The frequency of these most energetic events of the entire Solar System follows the 11-year solar cycle.

At the peak of the solar activity cycle, several dangerously high-intensity class flares and CMEs may occur (on average, around 2-3 per day at maximum). When a CME associated with a flare arrives to Earth several days later, its interaction with Earth's magnetosphere may trigger geomagnetic storms, with the additional (sometimes dramatic) consequences for various technological systems critical to our civilization (e.g., natural gas and oil pipelines, power lines, telecommunication, navigation etc). These societal assets and services are vital for the economic welfare and security of every citizen, but may be subject to failure due to solar activity, as it has indeed happened in the past (see e.g. [Eastwood et al., 2017](#)).

A crucial difference between today's world and the one from the many centuries ago is that we are now substantially more reliant upon computers and communications to run our commerce, work facilities, and even our forms of entertainment and recreation. The initial 15 geostationary and a few tens of Molniya-orbit communication satellites operated in 1989 have grown to more than several hundreds by now. Space weather may now affect such critical areas as: (1) the communication and navigation systems, (2) spacecraft operations, (3) global technological systems: power line distribution, pipeline operations, geophysical exploration, (4) research: balloon and rocket experiments, carbon dating, ozone measurements, and (5) the aircraft and commercial space travel (see, Figure 1 in [Pevtsov, 2017](#)). Our current economic development places even greater demands on the power network infrastructure and near Earth space-based assets. Due to such a reliance on global technological systems it is now believed that one of the biggest threat of civilisation breakdown may come from space weather. Therefore the question is not anymore "*whether*" it will occur, but "*when*" and "*how strong*". A direct consequence of and follow-up to this question is that one should improve a space weather early warning system by developing new capabilities for the observations and modelling of space weather precursors. This latter is the context of this current paper. With the Solar Activity Monitor Network (SAMNet) we now take a step in the direction to develop a ground-based facility for advanced space weather warning, in particular, to improve flare forecasting. The core scientific concept and underpinning scientific justification (see Section 2) behind SAMNet is the idea to exploit the line-of-sight (LoS) component **primarily** of the magnetic and **additionally** of the velocity (i.e. Doppler) fields observable not just at the visible surface of the Sun (i.e. in the photosphere) but at a range of heights in the lower solar atmosphere in order to improve the lead time for flare warning.

In this paper, we focus on the use of the measurable LoS component of magnetic data as this is a more popular and developed area for space weather pre-cursor applications. However, there is emerging literature about how the Doppler data of the near-side Sun may be used for imaging the emergence of active region of the far-side (see e.g. [Zhao et al., 2019](#)). Another growing application of the Doppler velocity fields is related to the evolution of magnetic helicity. In particular, the shearing and emerging components of helicity may provide additional and complementary pre-flare warning. How to determine these helicity components of the temporally varying magnetic helicity, how to assess their capability to serve as flare warning,

and how information on the LoS velocity components of the magnetised plasma will constrain the analysis, is demonstrated e.g. by [Ye et al. \(2018\)](#); [Korsós et al. \(2020b\)](#).

Novel flare prediction methods recently developed have already demonstrated, through observations and 3D simulations of Active Region (AR) dynamics, that information about the magnetic field *at various heights in the lower solar atmosphere*, in particular in the chromosphere, has the potential to considerably improve forecasting lead times. Therefore, a facility is now needed to provide a routine and continuous supply of magnetic and velocity field data of the entire solar disk, not just ARs, at *various atmospheric heights* and in particular the chromosphere. One needs to emphasise at this point that SAMNet and its core Solar Activity Magnetic Monitor (SAMM) instrument will be capable of acquiring the full disk magnetograms, which then can be used for a more in depth analysis of sub-areas around ARs.

After a brief summary in Sec. 2.1, the so-called weighted magnetic gradient (WG_M) method is introduced. WG_M is particularly developed to exploit measurements of a SAMNet-type facility. The method is based on assessing the evolution of the horizontal gradient of LoS component of magnetic fields observable by a series of SAMMs, the core instruments of SAMNet, in focused areas of ARs. Although we demonstrate the underlying science for ARs only, one has to recognise the importance of full disk magnetic observations. For example, CMEs often originate from (and their magnetic field is defined by) the chromospheric filament eruptions. Observing only ARs may miss other source regions of CMEs. Also, CME triggers may be remote, and so, without a global (full disk) view they could be missed too. In Sec. 2.2, arguments are also made using machine learning in favour of why a cost-effective set of low-resolution diffraction-limited ground-based solar telescope system like SAMNet may provide data with sufficient resolution and accuracy for flare eruption warning. In Sec. 2.3, it is demonstrated what gain in forecasting lead time may be achieved by applying the WG_M method to SAMM-type LoS magnetic data of the lower solar atmosphere. Here, the concept of optimum height is introduced where such LoS data of ARs may yield the best lead time (see Sec. 2.3.3). Next, in Sec. 3, the concept of MOF is introduced with applications to observe the lower solar atmosphere, followed by Sec. 4 devoted to the novel idea of tuning MOFs. Sec. 5 embarks on the introduction of the SAMM, an instrument realisation of mounting MOF-based filters on solar telescopes. The Solar Activity Monitor Network as an ensemble of SAMM solar telescope stations is introduced and discussed in Sec. 6. Finally, Sec. 7 discusses the potential future directions and limitations of a SAMNet-type of flare forecasting system.

2. Forecasting using LoS component of solar magnetic fields: WG_M method and ML approach as underpinning science

Here, we embark on a brief but necessary scientific justification in favour of the WG_M method that enables e.g. flare forecasting at source, using LoS solar atmospheric magnetic data that are obtained by e.g. a series of SAMM sentinels of SAMNet. In the Introduction, it was mentioned that we employ ARs as examples. However, the method may work equally well for the entire solar disk, subject to some limiting selection criteria outlined below (e.g. for reducing inaccuracy caused by measuring the magnetic field close to limb, etc.). It is also not claimed that the WG_M method is superior when compared to other methods. There are a number of excellent methods available in the literature that rely on a range of predictive parameters of solar eruptions (see e.g. [Georgoulis,](#)

2013; Barnes et al., 2016; Leka et al., 2018, 2019a,b, and references therein). However, most flare and CME forecast methods apply only photospheric magnetic and Doppler data of ARs for forecasting. Some recent, pioneering approaches with varying degrees of success attempt already to incorporate solar atmospheric extreme ultraviolet (EUV) data and/or use Machine Learning in order to improve forecasting accuracy (see e.g. Qahwaji and Colak, 2007; Bobra and Couvidat, 2015; Anastasiadis et al., 2017; Florios et al., 2018; Kim et al., 2019; Wang et al., 2019; Campi et al., 2019; Camporeale, 2019). Caveats of applying ML techniques is also addressed in Liu et al. (2021).

Detailed information on measuring, and the consequent modelling, of the 3D magnetic field structure of the solar magnetic field, in particular for an AR, would be important to obtain more accurate insight into the pre-flare evolution locally in the solar atmosphere. However, direct routine observations of the 3D magnetic field in the lower solar atmosphere, above the photosphere up into the top of the chromosphere, are currently not available. **An overwhelming majority of observations these days refer to either the LoS component or the full magnetic field vector in the photosphere only.** Nowadays, approximate methods for modelling the local magnetic field vector in the solar atmosphere include its construction using current free (potential, PF) or nonlinear force-free field (NLFFF) extrapolation techniques. In practice, however, to construct an accurate and reliable 3D magnetic field structure, also called the magnetic skeleton, of an AR from photospheric measurements is still a challenging task with a number of caveats, see e.g. Wiegelmann and Sakurai (2012).

The focus of this section is to demonstrate why it is sufficient for an efficient and accurate flare forecasting to observe (i) *only* the line-of-sight (LoS) component of the solar magnetic field B_{LoS} , and (ii) B_{LoS} at a *range of heights* in the lower solar atmosphere measured e.g. by an instrument suit like SAMMs of SAMNet. In Sec. 2.1, the core principles and applicability of the WG_M are outlined, while the section on employing machine learning (Sec. 2.2) is a key aspect to demonstrate that the skill score is basically not changing drastically by binning the HMI data. This latter information is in support of applying low-resolution, basically seeing-limited (and also diffraction limited) LoS magnetic data.

2.1. The WG_M method

It is shown by e.g. Schrijver (2007) that the most intense flares are in connection with the location of the strongest magnetic gradient. The *weighted horizontal magnetic gradient* (WG_M) method was developed and introduced originally by Korsós et al. (2015, 2019, 2020a) for the pre-flare tracking of ARs in the photosphere. The concept of the method is based on the evolution of the weighted horizontal magnetic gradient of the LoS component of the magnetic field (WG_M) between two opposite polarities in a δ -spot at the solar surface. Therefore, one needs to identify the δ -spot(s) of the selected ARs for the WG_M method. To find a δ -spot, we adopted and apply the automatic PIL recognition program developed by Cui et al. (2006).

Figure 1a illustrates the methodological framework of the WG_M method applied to an arbitrary example, say AR11166. The WG_M quantity itself is defined as a proxy to measure the magnetic non-potentiality in a contoured-up region of interest where D_{pn} is the distance between the area-weighted centers of the positive and negative polarities (barycenters) in this cluster, see Fig. 1b.




Fig. 1. Diagnostic diagrams of a sample investigated AR 11166. The upper panel of column a) is the variation of the WG_M parameter and the lower panel shows the evolution of the distance (D_{pn}). The right panels in column (b) are the white light (upper panel) and LoS magnetogram (lower panel) images of AR 11166 taken at 07:59 on 9 March 2011.

AR 11166 produced a strong X1-class flare according to the GOES flare classification system. The pre-flare dynamics and the related physical processes at the solar surface were investigated using data with an hourly temporal resolution from joint HMIDD ground- (Debrecen Heliophysical Observatory, DHO) and space-based (Helioseismic and Magnetic Imager on-board Solar Dynamics Observatory, SDO/HMI) sunspot data catalogues (see, e.g. Baranyi et al., 2016) from 2010 to 2014. The WG_M analyses focused on the vicinity of polarity inversion line(s) (PILs), which is within the red ellipse(s) in Fig. 1b. This is the area where the indicative characteristic features of the imminent flaring behaviour appear up to two-three days prior to the actual flare onset.

By inspecting Fig. 1a, two important findings of the WG_M method are as follows: i) First, a prominent and characteristic pre-flare patterns of the WG_M proxy quantity is identified: increasing phase, maximum and gradual decrease prior to flaring. The overall pre-flare behaviour of the WG_M parameter is approximated by aqua "inverted V-shape" on the top panel of Fig. 1a. This characteristic pre-flare behaviour is also confirmed by Korsós et al. (2015, 2019) in further approximately 130 flare cases between 1996-2014, where a linear relationship was found between the pre-flare WG_M maximum (WG_M^{max}) and the largest flare intensity class of the AR investigated. This relationship yields a tool to *estimate the expected flare intensity* following the preceding maximum of WG_M . ii) Further, the introduction of WG_M enables to determine a second flare precursor. Namely, the two barycenters of opposite polarities display a unique characteristic pattern of converging and diverging motions prior to the flare, which is illustrated with a red "U"-shaped parabolic curve in the lower panel of Fig. 1a. This "U"-shaped precursor allows to *estimate the expected onset time of flare* in a δ -type AR.

In summary, applying the WG_M -method to photospheric data only yields an about 7 hrs gain in lead time. Here, at this point, we wish to emphasise that achieving this progress is made by employing a relatively low spatial resolution (of the order of an arcsec) that is typical for ground-based observatories, as **would be expected from e.g. SAMNet as a lower limit**, routinely monitoring the solar surface with a cadence of **as low as** about an hour. **SAMNet will have, however, a better cadence at about 10 mins for a fixed wavelength that enables then a more accurate flare warning.**

2.2. Application of ML to LoS magnetic fields

In this section, we support the efforts that rely on how robust space weather forecasting, using LoS magnetic data, can be against resolution issues, i.e. how suitable a **ground-based** seeing-limited instrumentation is. We show that downgrading the HMI data does not really influence the skills cores/probabilities of forecasting.

ML methods can be used to build a solar flare forecasting model (Camporeale, 2019). Some of these rely on parameters calculated from the observational images (Nishizuka et al., 2018; Wang et al., 2020); others directly input solar images and automatically extract forecasting patterns (Huang et al., 2018; Park et al., 2018). We apply convolutional neural networks (CNN), which can process images directly, to explore the influence of image resolution on the performance of solar flare forecasting model. CNN is mainly composed of the convolutional layer, the nonlinear layer, the pooling layer and the fully connected layer. The convolutional layer is applied to extract forecasting patterns, the nonlinear layer provides the nonlinear transform in the forecasting model, the pooling layer is used to reduce the complexity of the forecasting model, and the fully connected layer is the last layer in CNN, the relationships between forecasting patterns and solar flares are obtained in this layer.




Fig. 2. Flow chart for exploring influence of resolution on model performance.

The original HMI magnetograms of active regions and their downgrading images are fed into the CNNs, respectively, as shown in Fig. 2. The models can provide forecasting results by us-

ing magnetograms with different resolutions. By comparing the performances of these models, we can estimate the influence of image resolution on forecasting model. Using bi-cubic interpolation method, we now downgrade the original image $1/\alpha$ ($\alpha = 10, 20, \dots, 100$) times. The flaring sample is considered as positive sample, meanwhile, the non-flaring sample is considered as negative sample. Hence, there are four possible outcomes in the solar flare forecasting model: i) True positive (TP) are that samples are correctly predicted as positive; ii) True negative (TN) are that samples are correctly predicted as negative; iii) False positive (FP) are that samples are wrongly predicted as positive; and iv) False negative (FN) are that samples are wrongly predicted as negative.

Based on these possible outcomes, the true skill score (TSS) (Eq 1) is considered as the index to evaluate the model performance.

$$TSS = \frac{TP}{TP + FN} + \frac{TN}{TN + FP} - 1, \quad (1)$$

The performance of forecasting models built on magnetograms with different spatial resolutions is shown in Fig. 3. The result indicates that the ML-based forecasting model is not sensitive for the spatial resolution of the input images. The ML-based model can work well, even if the input image is downgraded to 1/90 of the original size for HMI magnetograms. This result is rather important for facilities like SAMNet that may not achieve very high resolution due to the limiting effects of seeing (and diffraction, unless going for large apertures, say above 30 cm or even larger).




Fig. 3. Performance of forecasting models trained on magnetograms with different spatial resolutions. Original images are downgraded $1/\alpha$ times.

2.3. The power of WG_M - Forecasting as function of height with SAMNet-type data

Let us apply the WG_M method to LoS magnetic data in the lower solar atmosphere to establish how to further increase the flare prediction capability of the method. We now describe and demonstrate with a typical example how the forecast lead time gained by applying the WG_M -method to photospheric magnetic data can be even further improved by applying it to 3D LoS magnetic data at a range of heights across the lower solar atmosphere that may be delivered by a SAMM instrument.

For the demonstrating the power of applying LoS magnetic data at a range of heights, a 3D magnetic map data catalogue is constructed, which includes the area, mean magnetic field data and the location (Carrington coordinates, L for longitude and B for latitude) of sunspots of ARs at every 45 km in height from the photosphere into the lower corona on an hourly bases for the selected ARs. Eligible ARs now have to satisfy the a suit of selection criteria, see Korsós et al. (2019, 2020a). These criteria are a little different, i.e. more restrictive, from those when compared to the criteria of the photospheric event selection, applied in Sec. 2.1.

Once the above selection criteria are satisfied for an AR, first we use extrapolation to construct the 3D magnetic skeleton of this AR. Next, as a function of height, the WG_M -method is employed and the gain in lead time of flare onset forecasting is determined and compared to the counterpart lead time using photospheric data only. Finally, we search for a range of heights where best lead time improvement can be achieved. Such height will be called the *optimum height*. Once a solar magnetic monitoring system, e.g. SAMNet, would observe the LoS component of the magnetic field at the optimum heights in the lower solar atmosphere, this would then yield the best possible flare onset forecast.

2.3.1. 3D magnetic skeleton of Active Regions

It has recently been shown by Korsós et al. (2020a) that potential field (PF) extrapolation approaches are sufficient and there is no need for CPU/GPU-hungry and also time-consuming more complex extrapolations. For a demonstration, here, we use the HMI LoS magnetograms as a photospheric lower boundary for constructing the magnetic skeleton of two random examples, say ARs 11166 and 12192, each embedded in a 3D box (see Fig. 4). The ARs are tracked by using the




Fig. 4. The figure illustrates the three-dimensional magnetic field of a) AR AR11166 (left) and b) 12192 (right). The red-blue colour bar refers to the positive-negative polarity vertical B_z magnetic field values measured at the solar surface and computed as function of heights in the solar atmosphere. **Particular heights outlining approximately the heights of the line formation of K I, Na D, Ca I and He I are highlighted.**

Yet Another Feature Tracking Algorithm¹ (YAFTA; Welsch and Longcope, 2003; DeForest et al., 2007). For the PF extrapolation, we use the method published by Gary (1989). To extrapolate the

¹ YAFTA is available from <http://solarmuri.ssl.berkeley.edu/?welsch/public/software/YAFTA>

PF magnetic fields at every moment of time, we employed the LFFF IDL extrapolation code from www.heliodocs.com², which is based on Gary (1989).

Figure 4 illustrates the three-dimensional magnetic field of AR 11166 (left) and AR 12192 (right), respectively, using the PF method described above. **At horizontal crosscuts corresponding to the approximate heights of the line formation of K I, Na D, Ca I and He I chromospheric lines the LoS component of the magnetic field could be observed with a solar telescope (e.g. a SAMM) that is equipped with e.g. K I, Na D, Ca I and He I filters. The horizontal cuts at the formation height of the He 10830 Å line would be a height of "sunspot tracker" at the top of the chromosphere.** In the next section, the WG_M -method will be applied and evaluated to determine the lead time for flare forecast at cross-cut from the photosphere into the deep chromosphere to demonstrate the power of concept of SAMM-based SAMNet.

2.3.2. Application of WG_M to 3D magnetic skeleton of Active Regions

Korsós et al. (2018) identified the so-called optimum heights above the photosphere where the flare precursors of the WG_M method appear earliest prior to flares of a simulated AR. This finding has led Korsós et al. (2020a) to test and validate the modelling with real solar active regions. They embarked on analysing the variation of the flare onset lead time for 20 flares taking place in 13 ARs. They found that the lead time estimate can be improved by an additional 2-8 hrs when compared to the counterpart lead times obtained **only** from photospheric LoS magnetic data of the same ARs.

Let us put this in the context of yet missing routine chromospheric magnetic field observations where this gap will be closed up by SAMNet. Figure 5 shows the application of the WG_M method to the 3D magnetic field of ARs 11166 (left panel) and 12192 (right panel), respectively. The evolution of the value of WG_M and the distance parameter D_{pm} are plotted at particular heights representing the photosphere (upper left pair) and the line formation heights of K I (upper right pair), Na D (lower left pair) and Ca I (lower right). The x -axis is time, measured in days. For a given pair of panels, the upper panel is the temporal variation of WG_M . The pre-flare behavior of WG_M is fitted by an n th-order polynomial (red line), where the orange dot corresponds to the maximum of WG_M . The bottom panel demonstrates the evolution of D_{pm} . The consecutive maximum-minimum-maximum (orange-blue-orange dots) locations of the fitted n th-degree polynomial highlight the full converging-diverging motion uncovered by D_{pm} . **For details about how the minimum value of D_{pm} is determined, see e.g. Korsós et al. (2019).** The vertical green stripe indicates the moment when the flare occurred. Even a simple visual inspection shows that the characteristic pre-flare features vary as a function of height. The heights are chosen to match the approximate formation heights of the three lines (K I, Na D, Ca I) as these are popular lines to measure the chromospheric magnetic field.

Next, we demonstrate the role of optimum height, i.e. the height for achieving the longest gain in the lead time for estimating the flare onset using the WG_M method, and whether this height can be find by using a SAMM sentinel equipped with a series of filters (e.g. K I, Na D and Ca I, and He I 1083 at a later stage) to measure the required LoS magnetic fields in the lower solar atmosphere.

² http://www.heliodocs.com/php/xdoc_print.php?file=SSW/packages/spvm/idl/bff/bff.pro




Fig. 5. The figure illustrates the application of the WG_M method to the 3D magnetic field of ARs 11166 (left panel) and 12192 (right panel), respectively. The evolution of the value of WG_M and the distance parameter D_{pm} are plotted at particular heights representing the line formation heights of K I, Na D and Ca I. The green strip denotes the onset of the X2.0 flare observed. The vertical blue line marks the moment of closest position of the positive/negative barycentres at a range of heights including the photosphere and the line formation heights of K I, Na D and Ca I lines.

2.3.3. Optimum height for forecasting

The evolution of the WG_M and the D_{pn} are different at various heights, as has been described above (see Fig. 5). In order to optimise the flare precursor capability of the WG_M method, let us now identify optimum height(s) in the solar atmosphere. There, the precursor behaviours of the WG_M and D_{pn} parameters are identifiable earliest prior to each flare. The optimum height(s) are also where these parameters would concurrently yield the earliest signs of pre-flare behavior in time.

First, in Fig. 6a,b we plot the variation of the start time of the approaching phase (labelled as "Starting"), the moment of the closest approach (labelled as "Minimum") and the flare onset time of the X1.5 (left panel) and X2.0 (right panel) flares (red line) as a function of height. In Fig. 6, the filled grey squares mark the estimated line formation heights for the K I, Na D, Ca I and He I 1083 magnetically sensitive spectral lines. Most noticeable is that, in general, there are certain heights above the photosphere and well in the chromosphere where the approaching motions began hours earlier and reach the closest point of approach also hours earlier than at the photosphere or at other heights in the solar atmosphere. The actual numerical evaluation shows that the gained lead time is about 3/8 hours, would a flare prediction been attempted using chromospheric data, for the X1.5/X2.0 flare, respectively, when compared to their respective photospheric counterparts. This would be a definite (for the X1.5) or an even more considerable (for the X2.0) improvement in determining the flare onset time, would a solar instrumentation measuring routinely LoS magnetic data in the chromosphere. A more comprehensive analysis of [Korsós et al. \(2020a\)](#) has shown that, in spite of the apparent limitations of employing PF extrapolations only, the potential lead-time improvement for estimating the flare onset time varies in the interval (2, 8) hours if one applies the WG_M method to an identified "sunspot" in the height region between ~ 800 - 1000 and ~ 1800 km




Fig. 6. The figure illustrates the application of the WGM method to the LoS component of the 3D magnetic field skeleton of a) AR 11166 (left) and b, AR 12192 (right), respectively, to determine optimum forecasting height. The moments of starting and closest convergence (i.e. minimum) times are deduced by the maximum and minimum values, respectively, of the best n th degree polynomial fit to the D_{pn} data. The colour bar gives information about the actual value of D_{pn} . **The grey zones are only indicative line formation heights.**

above the photosphere.

Let us now summarise the main science pillars and findings of Sec. 2 that supports the concept of constructing SAMMs with the purpose of improving flare forecasting. It is shown that it is sufficient for yielding an efficient and accurate flare forecasting to analyse *only* the line-of-sight component of the solar magnetic field B_{LoS} , and that analysing B_{LoS} at a *range of heights* in the lower solar atmosphere opens up to gain a considerable forecasting lead time. No high resolution is necessary, seeing or diffraction limited observations would be sufficient. Now, in the following sections, we outline how to realise these science pillars in practice with a facility that will continuously monitor the solar atmospheric magnetic field. This will be done by employing specific magneto-optical filters on solar telescopes that have the task to continuously and routinely monitor the lower solar atmosphere.

3. The magneto-optical filter (MOF)

This section is addressing the concept of MOF itself explaining the underlying basic idea and frequency transmission (Sec 3.1) while also showing some trial images (Sec 3.2). **Given the new realisation context and application of MOF technology to space weather, a brief overview is given in order to better understand the limitations and what future options for development may lay ahead.**

3.1. The MOF concept

The magneto-optical filter (MOF) is a frequency-selecting element, which utilises the Faraday effect: the plane of polarisation of linearly polarised light rotates as it propagates through a medium that has an axial magnetic field applied. Linearly polarised light can be thought of as being composed of equal amplitudes of left- (LCP) and right-hand circularly polarised light (RCP). In a circular birefringent medium where the refractive indices for LCP and RCP are different the two components experience different phase shifts, resulting in a net rotation of the linear polarisation. The amount of optical rotation strongly dependent on the dispersion of the medium. When an atomic vapour is used as the dispersive medium, the Zeeman effect splits the atomic absorption lines into two components, σ_+ and σ_- transitions. Hence, the optical rotation only occurs near atomic resonances, which are intrinsically very narrow. This makes MOFs ideal instruments for filtering out unwanted optical background noise. It is for this reason that the MOF is employed across many disciplines and in solar physics it referred to as Cacciani Filter after [Cimino et al. \(1968\)](#), [Cacciani and Moretti \(1994\)](#), [Cacciani et al. \(1997a,b\)](#). In addition to solar physics, magneto-optic filters also find a wide range of applications in other disciplines. Examples include e.g. filtering of frequency-degenerate photon pairs ([Zielińska et al., 2014](#)); quantum key distribution ([Shan et al., 2006](#)), laser-frequency stabilisation ([Miao et al., 2011](#); [Keaveney et al., 2016](#); [Chang et al., 2019](#)); Doppler velocimetry ([Bloom et al., 1993](#)); atmospheric LIDAR ([Yang et al., 2011](#); [Xia et al., 2020](#)); and simultaneous atmospheric wind and temperature measurement ([Huang et al., 2009](#)).




Fig. 7. A schematic illustration of the complete MOF instrument for photospheric observations. Collimated sunlight passes through a two-stage filtering process. The filter section consists of an atomic vapour subject to an axial magnetic field placed between crossed polarisers. The temperature of the vapour and the strength of the magnetic field are chosen to achieve a suitable narrow transmission profile – with two wings – centred on the atomic resonance. The second vapour cell and quarter waveplate are used to encode the transmission of the opposite handedness of circularly polarised light into vertical and horizontally polarised light, which are subsequently imaged by cameras CMOS1 and 2, respectively.

Figure 7 illustrates the principle of using the complete MOF instrument necessary for photospheric observations, which consists of a filter section MOF and a wing selector MOF. Examples of the transmission profiles through a filter section sodium MOF, for one particular input polarisation, are shown in Fig. 8. A wavelength of 0 mÅ refers to the unperturbed sodium transition ($3^2S_{1/2} \rightarrow 3^2P_{3/2}$), i.e. atomic resonance. Typically, the spectrum has two wings, in Fig. 8 (b), split

from the atomic resonance by the Zeeman effect. The wing blue-detuned from the atomic resonance, i.e. the wing on the left of 0 mÅ in Fig. 8 (b), is referred to as the blue wing (B) and the wing on the right of 0 mÅ is known as the red wing (R).




Fig. 8. The simulation and experiment results of the sodium MOF transmission functions. (a) and (b) show the transmission functions of Na D MOF with a center peak and with two wing peaks, respectively. Panel (c) shows the sodium MOF filter that has a single peak in the Fraunhofer D2 line center, enabling chromospheric filtering. The red solid line is the Fraunhofer line and the blue and black solid line are the MOF transmission functions.

Depending on the application, the transmission through the filter section MOF can be tailored to give desirable features. For solar chromospheric observations, a transmission that has a single peak centered on the atomic transition is required (Fig. 8 (a)), whereas, for both Doppler- and magnetic-field observations, a transmission that has two wings is necessary (Fig. 8 (b)). The transmission shape and passband can be tailored by choosing appropriate filter section MOF parameters - software has been developed by Yang et al. (2011) to optimize these parameters, which agree with experimental transmission profiles as shown in Fig. 8.

For photospheric observations, where a filter section MOF transmission with two wings is necessary, a second MOF is required to extinguish the unwanted wing. This is referred to as the “Wing Selector MOF” in Fig. 7. The light that exits the filter section MOF, and subsequently enters the wing selector MOF, can be decomposed into RCP and LCP. For the red wing, the LCP component is absorbed by the wing selector MOF, whereas for the blue wing, the RCP component is absorbed. Using a quarter waveplate and polarising beam splitter (PBS) after the wing selector enables the LCP and RCP components to be separated, such that light in the blue wing is detected on CMOS1 (B) and light in the red wing is detected on CMOS2 (R). Since the input polarisation to the filter section MOF can be set to address either σ_+ or σ_- transitions, there are a total of four images that can be taken: R^+ and B^+ for σ_+ input and; R^- and B^- for σ_- input.

Using these four images, the velocity v_{LoS} and magnetic field B_{LoS} in line-of-sight can be derived:

$$B_{LoS} \propto \frac{R^+ - B^+}{R^+ + B^+} - \frac{R^- - B^-}{R^- + B^-}, \text{ and } v_{LoS} \propto \frac{R^+ - B^+}{R^+ + B^+} + \frac{R^- - B^-}{R^- + B^-}. \quad (2)$$

Measuring v_{LoS} and B_{LoS} serve as strong boundary conditions for modelling solar magnetic fields, particularly in the lower region of the solar atmosphere where the magnetic field is highly non-force-free.

3.2. Application of MOF to observing the solar atmosphere

The MOF cell was first envisaged and used for solar applications, in particular in the context of observations of the lower solar atmosphere, by Cimino et al. (1968, 1970); Cacciani et al. (1990, 1997a,b). Full disk maps of the Doppler velocity and magnetic field strength at the photosphere can be obtained through the use of the vapour cells coupled with polarisation active electro-optical components (see, e.g. Oliviero et al., 2002; Finsterle et al., 2004b; Murphy et al., 2005; Moretti et al., 2010; Forte et al., 2017, 2018).

The formation height of the line center of the Fraunhofer Na D2 is in the chromosphere and those of the line wings are in the photosphere (Kuridze et al., 2016). The single peak MOF selected the line center and blocked most of the line wings of the Fraunhofer Na D2, therefore the chromosphere could be observed. A sodium MOF with a single peak and the sodium MOF




Fig. 9. Observing the lower solar atmosphere. The x - and y -axis represent the position in pixels. One pixel corresponds to 0.3 Mm. The resolution of the z -axis is one tenth of the resolution of x - and y -axis. a) Left is the visualisation of the lower solar atmosphere embracing NOAA AR 11745 (upper left sunspot group) and NOAA AR 11744 (sunspot group in the middle-right). The bottom panel is the photospheric magnetogram, the top panels are taken by the sodium MOF with two peaks (600 - 700 km) and single peak (800 - 850 km) at UT20130513T030854. Because the vertical scale is much larger than the horizontal one, the bottom panel is hardly visible. b) Right is the view from the top.

with two peaks were employed to observe the Sun. Fig. 9a shows a test magnetic skeleton of two ARs, namely ARs 11745 (upper left sunspot group) and NOAA AR 11744 (sunspot group in the middle-right) image of the lower solar atmosphere. The test observation is to demonstrate the application of a MOF-based approach of observing the lower part of the solar atmosphere with the two peaks MOF. Underneath the MOF image is the corresponding HMI magnetogram. The chromosphere was observed with the single peak MOF as seen in Fig. 9b, top view. This example demonstrates how the LoS component of the magnetic field can be obtained and constrained by observations in the most critical (optimum) height of the lower solar atmosphere. Taking such

images at a cadence, say 10 mins **at fixed wavelengths**, would enable to apply the WG_M (or any other similar) method for flare warning.

Here, we only show an example of the use of the Na D line (Uitenbroek, 2006; Kuridze et al., 2016) with a MOF-based solar telescope. However, other lines (e.g., K I, Ca I or even He 10830 Å) with different formation heights (see e.g. for K I Quintero Noda et al. (2017); Ca I Supriya et al. (2014) and for He I Avrett et al. (1994); Leenaarts et al. (2016)) would provide B_{LoS} at further heights, yielding additional constraints on limiting the freedom of constructing the magnetic skeleton in the lower solar atmosphere even with PF method. Once such a skeleton is obtained, given that the PF methods are executable almost instantaneously, even on a modest laptop, approximately near real-time flare warning could be issued. This is the true strength of a MOF-based solar telescope, the core to SAMNet.

4. The tuneable magneto-optical filter (T-MOF)

Here, we now briefly discuss the novel idea of the tuneability of a MOF by changing its cell temperatures. When the tuneability capability of MOFs is fully developed, this will have important implications in space weather, solar and astrophysics applications. An initial application to solar physics is shown via a series of intensity images as a result (see Fig.10) and an associated example of simulations of a sample sodium MOF line width with the demonstrated effect of tuneable filtering (see Fig.11). The tuneability of MOF is a yet uncharted approach, while being a competitive alternative to other, well-established measuring techniques of the solar magnetic field, e.g. by Fabry-Perot or Lyot filters.




Fig. 10. Observing the lower solar atmosphere by a tuneable sodium MOF application on 30 July 2013, showing the solar intensity. The temperature of the MOF varied from 160° C to 220°C. UTC time and temperature values for each panel are labeled on the snapshots individually. **The sequence of images outline an intensity map scan descending from the mid-chromosphere downward toward the photosphere.**

During our solar observations at the SAMNet Wuhan Solar Observing Station on 30 July 2013, the variation of temperature of the MOF cell during the observations was tested. The **initial** temperature of the MOF was varied from about 160°C to 220°C. **At the same time, the passbands of the wing-selector MOF may also need to be tuned (by changing the magnetic field for the wing selector).** The associated full disk snapshots are shown in Fig.10. The snapshot in the left top is for the initial, lowest temperature of the MOF, and snapshot in the bottom right is for the highest one. The temperature and UTC time are labeled for each snapshots. The initial chromospheric features are shown in the top left (Fig.10a). By varying the temperature, one scans through the line and arrives at observing the solar photospheric features as seen in the snapshot in bottom right (Fig.10h). Given the behaviour of the Na D line (see Fig.11 or Yang et al., 2011), each temperature




Fig. 11. Simulation results of transmission function for the sodium MOF. The temperature of MOF is changed from 160°C to 220°C. The magnetic field is 200 mT, and the length of cell is 25 mm. The Na D₂ line is from (Wallace et al., 2011).

of the MOF maps a relatively narrow band of the solar atmosphere. **We note that with the temperature tuning capability the cadence is slightly varying when compared to observations at a fixed wavelength as time is now needed for the vapour to settle to its newly adjusted temperature.** However, on average, a line -and therefore the lower solar atmosphere- can **still** be scanned in about 30 mins **depending on the number of line positions.** This is a **promising initial** result and well suited for further constraining the much-needed magnetic skeleton of the lower solar atmosphere as even just intensity contours may serve to be a guidance for the computation of the local magnetic field strength.

In summary, the transmission function of a given MOF with fixed atomic cell could then be "tuned" by changing its temperature (or, for the record, even the magnetic field) and observe differ-

ent heights of the Sun's atmosphere. Note that this is equally true, though not demonstrated here, to other suitable MOF cells, based on other atomic vapours (e.g. K I, Ca I and He I).

5. Realisation of MOF-based telescope for solar physics

5.1. Solar Activity Magnetic Monitor - SAMM

Here, after a short historic account, we briefly outline the current realisation of a MOF-based solar telescope that has been built on these earlier developments. This new design and developments is the so-called Solar Atmospheric Magnetic Monitor (SAMM) station device, that allows to observing routinely the solar B_{LoS} at various heights (e.g., K I and Na D). In this section, we also describe very briefly the optical telescope assemblies (OTAs), cameras, etc.

5.1.1. Heritage Context

[Agnelli et al. \(1975, 1976\)](#) introduced preliminary observations by the MOF Na D lines and obtained one of the first images of the Sun by this technology on 11 May 1974 at UT11:40 (see their Fig 5a.). The specific idea of measuring lower solar atmospheric velocities was further explored by [Cacciani and Fofi \(1978\)](#) and it was applied to measuring the Doppler signatures of the global solar acoustic (p) modes (e.g. [Rhodes et al., 1984](#); [Tomczyk et al., 1993, 1995](#)). In these papers, the practical realisation of the working principles of a MOF-based instrument, earlier only shown in theory by e.g. [Cimino et al. \(1968, 1970\)](#), [Beckers \(1970\)](#), [Beckers and Wagner \(1970\)](#) and [Cacciani et al. \(1971\)](#), was described in details with the associated transmission profiles. These pioneering studies were among the first to actually demonstrate how the underlying MOF-based concept can be put for real solar observations.

VAMOS (Velocity And Magnetic Observations of the Sun) - a solar atmospheric MOF imager, based on [Cacciani and Fofi \(1978\)](#); [Moretti et al. \(1997\)](#), has emerged some decades later. The first version of this instrument ([Cacciani et al., 1997a](#); [Moretti et al., 1997](#)) with two sodium vapour cells observed in the Na I D lines ([Oliviero et al., 1998a,b](#)). A newer 40 cm aperture telescope with two potassium vapour cells became operational in May 1999 located at the Capodimonte Observatory (see [Severino et al. \(2001\)](#) or Fig. 12 in [Oliviero et al. \(2002\)](#)). VAMOS obtained high-cadence



Fig. 12. VAMOS at the Capodimonte Observatory, Italy. Adopted from [Oliviero et al. \(2002\)](#).

observations of the Sun's intensity, velocity and longitudinal magnetic field component at the photospheric level at the K I 769.9 nm line formation height. The instrument's main characteristics are listed in Table I of [Oliviero et al. \(2002\)](#).

MOTH-II (Magneto-Optical Filter at Two Heights) - One of the earliest manifestations of developing a MOF-based solar telescope was the MOTH Doppler-magnetograph, upgraded to MOTH-II with two identical instruments, except the actual MOF cells and the narrowband interference filters. The MOTH device is described in [Finsterle et al. \(2004a\)](#), and a good overview can also be found [here](#). The concept of MOTH-II is identical to prototype SAMM's: deliver with a pair of, in this case, 20 cm aperture OTAs measured full disk Dopplergrams and magnetograms at two heights in the lower solar atmosphere. The K I line is used for photospheric and Na D2 line for lower chromospheric measurements. About the data pipeline and the superb observational results see Forte et al. (in this volume) and [here](#). Excellent applications, e.g. three-dimensional topography of the solar chromospheric canopy (see [Finsterle et al., 2004b](#)), the observing of traveling acoustic waves in the lower solar atmosphere ([Haberreiter et al., 2007](#)), or even potential space-based applications ([Stangalini et al., 2011](#)) all demonstrate the power of acquiring the magnetised lower solar atmosphere by MOF-based solar instrumentation.

SAMM - The technical realisation led by Avalon Instruments, supported by an initial start-up incentive from the Ministero per lo Sviluppo Economico (Ministry for Economic Development, MiE), Italy and additional supports from the UK (via STFC, U of Sheffield and QUB) as well as from Hungary, the very first stage of a further proof-of-concept dual-channel robotic telescope has already been initiated, designed, built and commissioned with 2 OTAs (using K I and Na D MOFs, see for the technical details at [Stangalini et al. \(2018\)](#) and hspf.eu/filters.html). This 2xOTA solar telescope is a scaled-down version of the full SAMM concept (see images on the right of Table 1) with the aim of demonstrating synoptic data provision only at two heights in the lower regions of the chromosphere (see the top two rows left of Table 1 for the line, wavelength, formation height range and diffraction limit).


Line	Wavelength [nm]	Height km	Diffraction limit resolution [km]	
K I	770	3-400	0,71	
Na D ₁	590	6-700	0.54	
Ca I	422	1000	0.39	
He I	1083	1900	1.19	

Table 1. Left: Wavelengths and **indicative** formation heights of the K I, Na D₁, Ca I and He I 1083 chromospheric lines. Right: Snapshot of the scaled-down SAMM prototype telescope at various stages of its design and development.

The prototype scaled-down SAMM observatory is an enclosure measuring approximately 2m on each side. Given its size, there is an enabled flexibility of portability making it suitable of an easy deployment, although this latter feature is more a bonus as opposed to being an essential requirement. Irrespective of its size, SAMM is provided with a mini-weather station, including rain sensors and a built-in internal camera for a secure remote surveillance of operations. In case of loss

of power or unfavourable weather conditions automated closure of the dome will take place within about a minute thanks to a backup battery and UPS that can provide operation for up to 20 minutes. SAMM's own computer unit provides control of both the sentinel and the instrumentation as well. The computer system is a standard PC, currently an Intel Xeon X5560 with dual CPUs at 2.80 GHz for a total of 8 cores, with 24 GB of RAM installed (see [Stangalini et al., 2018](#)). For the full-scale SAMM the computer system needs to be updated e.g. with Intel NUCs or similar.

This robotic test instrument, when fully operational, is hosted at Gyula Bay Zoltan Solar Observatory ([GSO](#)), the scientific and operational HQ of SAMNet, thanks to an open-ended long-term agreement of in-kind support secured from the Local Govt. at Gyula (Hungary). The robotic dome, now capable of hosting the 2-channel scaled-down version of the current version of the SAMM unit master station and associated initial data processing facilities (currently 64 TB) is under calibration testing at its developer's site (DS Group Headquarters, Aprilia, Rome, Italy).



Fig. 13. SAMM at DS Group HQ, Aprilia, Rome, Italy in May 2017.

5.2. Expanding the capabilities of SAMM

Currently, there are two potential developments to expand the monitoring capabilities of a SAMM station in order to survey the magnetic field and large-scale dynamics in the solar atmosphere. An option is to add another OTAs equipped with Ca I or He 1083 MOF capability for measuring the LoS component of the chromospheric magnetic field about one Mm (Ca I) and 1.9 Mm (He 1083) above the solar photosphere. The He MOF is rather challenging (see [Murphy et al., 2005](#)) but the Ca I MOF for SAMNet is under development (see Sec. 5.2.1). Another option is to mount coronagraphy capabilities as a 3rd or 4th OTA, based on the heritage and design of the High-resolution Spectrometer for Coronal Emission Lines (HISCEL) instrument with polarimetry capability. The HISCEL development is led by Aberystwyth University, and up-to date information is [here](#). This would enable obtaining imaging and velocity information above limb for CMEs lifted off the solar atmosphere.

5.2.1. The Ca I MOF cell

Potassium and sodium MOFs are routinely used in solar studies, however, to the best of our knowledge, there is currently limited application of the Ca-I MOF to solar physics. Although the Ca-I MOF is already commercially available (e.g. at [Mojave Solar](#)) and is occasionally




Fig. 14. Graphs showing computed properties of a calcium MOF utilising the $\text{Ca}(4p^1P_1 - 4s^1S_0)$ transition at 422 nm. The abscissa is detuning – difference between the frequency of the light and the atomic resonant frequency – in units of the Doppler width, $\Delta\nu_D$. An applied external magnetic field leads to a splitting of the upper state's energy level through the Zeeman effect. Graph a) shows the absorption profiles, α_+ and α_- , for the circular polarisations of the signal light, LHC that drives σ_+ transitions, and RHC that drives σ_- transitions, respectively. The magnetic field strength is 0.04 T, and the temperature $T = 720$ K. In b) the corresponding dispersion profiles are plotted for the σ_+ and σ_- transitions for a cell of length 50 mm. The Faraday rotation angle, $\delta(\nu)$, is plotted in c) for varying magnetic field strength. Note that optical rotations of the order, or greater than, π are generated for these parameters, as is necessary for efficient atomic filtering. In d) the transmission profile is calculated for each of the associated Faraday rotation angles in c). The characteristic two-wing spectral profile is evident.

used to observe the Sun, there is no routine synoptic observing implementation. The application of a Ca-I MOF to expand the surveying range of the lower solar atmosphere of a SAMM sentinel would be a real advantage, because the magnetically-sensitive Ca-I line has a formation height that is about one Mm above the solar surface (Supriya et al., 2014). Observations of B_{LoS} at these height ranges would further constrain the construction of the 3D magnetic skeletons of ARs, irrespective of whether there is a small overlap with the formation height of the Na D line. Therefore, this would have the additional benefit of improving our capability of delivering additional lead-time gains even further. Moreover, a calcium MOF allows observations of the

extremely interesting chromospheric layers of the solar atmosphere where the plasma-beta is approximately one. Ultimately, observing the three different magnetically-sensitive spectral lines (one photospheric and two chromospheric, see Table 1), it is clear that SAMM would produce superior, multi-height synoptic LoS magnetograms.

A calcium filter section MOF (see Fig. 7 for descriptor reference) was simulated utilising the Ca ($4p^1P_1 - 4s^1S_0$) strong transition at 422 nm, using an axial magnetic field strength of $B = 0.04$ T, a cell length of 50 mm and a vapour temperature of $T = 720$ K (Fig. 14). As outlined in section 3.1, the input polarisation of light, i.e. LCP or RCP, determines whether σ_+ or σ_- transitions are addressed, respectively (Adams and Hughes, 2019). Figure 14 a) shows the computed absorption coefficients, α_{\pm} , for LCP and RCP as a function of detuning (the difference between the frequency of the light and the atomic resonant frequency) in units of the Doppler width, $\Delta\nu_D$. Also shown are the corresponding refractive indices n_{\pm} (Fig. 14 (b)), whereby the differences in the absorption and refractive index curves, with respect to LCP and RCP, lead to optical rotation, $\delta(\nu)$, (Fig. 14 (b)) and filtering effects (Fig. 14 (d)), which are evidently dependent on magnetic field strength. The characteristic two-wing spectral profile is evident in (Fig. 14 (d)), which is necessary in photospheric studies. Further simulations have shown a sensitivity to the vapour temperature and, correspondingly, the number density of the atoms in the cell as well, since this is calculated as a function of temperature (Lide, 1995). Through use of computational optimisation procedures we can determine filter parameters that are optimum for our applications of interest (Keaveney et al., 2018).

6. SAMNet - Solar Activity Monitor Network

6.1. The SAMNet concept

In support of the successful latest development of applying MOF-based technology to solar observation, the Solar Activity Monitor Network was formed of individual identical Solar Activity Magnetic Monitor (SAMM) solar telescope stations at each partner site when at full capacity. Current international scientific partners with various levels of support and with principal interest include Kanzelhöhe Solar Observatory (Austria); NAOC Beijing, Huairou Solar Observatory, Mingantu Radio Observatory, USTC Hefei, Yunnan Astronomical Observatory, Xinjiang Astronomical Observatory and Innovation Academy for Precision Measurement Science and Technology (China); National U of Colombia, Bogotá (Colombia); Hvar Observatory (Croatia); Abastumani Astrophysical Observatory (Georgia); Eötvös U Budapest and the Hungarian Solar Physics Foundation (Hungary); IIA Bangalore (India); Catania Astrophysical Observatory, Rome Astronomical Observatory (Italy); Pulkovo Observatory, Kislovodsk (Russia); Astronomical Institute of the Slovak Academy of Sciences, Tatranská Lomnica (Slovakia); GONG site at Big Bear Solar Observatory (BBSO) and the Sunspot Solar Observatory of New Mexico State University (USA). The instrumentation suit may be supplied by Dal Sasso - Avalon srl (Rome, Italy) although alternative options are considered too. In order to achieve maximum performance without spending considerable effort on instrumental cross-calibration across the sentinel stations, it is a highly desired requirement -though not necessary- to have identically built SAMM stations at all partner sites. For the geographic distribution of the current SAMNet solar observatory partners see Fig. 15. A brief summary about the geographic location of the partner observatories

SAMNet team: Solar Activity Monitor Network

comprising SAMNet see Table 2. In addition, additional UK expertise (Durham University, QUB, U. of Aberystwyth, UCL/MSSL) with solar technology know-how have also joined the network.

Name	Abbrev	Lat	Long	Alt (m)	Instrument
Abastumani Astrophys. Obs.	AAO	N41° 45' 15"	E42° 49' 10"	1,650	530 mm Lyot cor 115 mm Lyot cor dual Ch-Ph telescope
Big Bear Solar Observatory	BBSO	N34.2614°	W116.9212°	2,000	GONG Network
Fuxian Solar Observatory	FSO	N24° 34' 48" N29° 09' 01"	E102° 57' 01" E100° 04' 13"	1,720 4,700	NVST Liu et al. (2014) CGST Deng (2011)
Huairou Solar Obs. Station	HSOS	N40.32°	E116.62°	64	35 cm SMFT; 60 cm StCT 10 cm Fe I vector m.f. 20 cm H-alpha full disk
Mingantu Observing Base	MOB	N42° 12'	E115° 15'	1,365	MUSER Yan et al. (2009, 2021) ISP Yan et al. (2018)
Nanshan Station	NS	N43.8°	E87.1°	2,080	H-alpha with Lyot filter
Wuhan Solar Obs. Station	WSOS	N30° 31' 54"	E114° 21' 26"	80	Na and K MOFs
Nat. Astron. Obs. Colombia	NAO	N4° 38' 23"	W74° 05' 00"	2,600	FiCoRI
Gyula Bay Zoltán Sol. Obs.	GSO	N46.649257°	E21.269346°	87	SAMM master SAMNet HQ
Hvar Observatory	HO	N43° 10' 38.14"	E16° 26' 56.29"	240	21.7/13 cm Ph/Ch-doublet
Indian Astron. Obs. (Leh)	IAO	N33° 43' 59"	E78° 53' 48"	4,500	NLST Hasan et al. (2010) H α Ravindra et al. (2018)
Catania Astrophys. Obs.	INAF-OACT	N37° 31' 43.71"	E15° 04' 17.38"	197	150-mm Cook ref. 150-mm H α Lyot Global High-Res. H α Network
Kanzelhöhe Observatory	KSO	N46° 40.7' N	E13° 54.1' E	1,526	Patrol telescope sys. H α , Ca II Global High-Res. H α Network
Kislovodsk Mountain Astron. Station	KMSA	N43° 44.77' N	E42° 31.42' E	2,070	53 and 20 cm coronagraph STOP telescope
Lomnický Stit Observatory	LSO	N49° 11' 42.53"	E20° 12' 47.43"	2,634	CoMP-S Kucera et al. (2010)
Sunspot Solar Observatory	SSO	N32.5°	W105.5°	2,807	Dunn Solar Telescope

Table 2. List of SAMNet partner observatories with their name, abbreviation, longitude, latitude, altitude and reference to solar instrumentation (if applicable). For FiCoRI see Guevara Gómez et al. (2017), for KSO see Pötzi et al. (2015, 2016, 2018); Veronig and Pötzi (2016) for the ESA Space Situational Awareness (SSA) programme. For KMAS see here, and for their observational data here.

SAMNet is now a UK-led international network with its scientific and observing HQ at Gyula Bay Zoltan Solar Observatory (GSO, see Tables 2 for some basic information) under the auspices of HSPF. Once at its peak performance, SAMNet will continuously monitor the Sun's magnetic and velocity fields both in the lower solar atmosphere (from photosphere to upper chromosphere) using a set of identical SAMM sentinel stations equipped with a range of magneto-optical filters (MOFs). Sentinels with suitable observing conditions will also have HISCEL-based set of coronagraph-mounted spectrograph as well imaging the low corona by means of a number of coronal visible lines in order to

- provide advanced warning of adverse space-weather events in the form of solar flares or coronal mass ejections (CMEs) that threaten our high-tech civilisation and the technosphere;
- provide advanced warning of the arrival of CMEs that are already formed and lifted off the solar surface; and

- provide much sought-after continuous synoptic maps (e.g., LoS magnetic and velocity fields, intensity) of the lower solar atmosphere.

The flare and CME onset warning is based on the WG_M method as outlined in Sec. 2, while the CME arrival will be predicted routinely using a new tool, CME Arrival Time Prediction using Machine Learning Algorithms (CAT-PUMA), developed by Liu et al. (2018). Even if only low corona imaging observations are available, prediction of CME arrival at L1 is still expected as demonstrated by e.g. Wang et al. (2019).

The Consortium of 23 partners from observatories, academia and industry is committed to investigate the feasibility, design and when funding is secured to develop its fully-robotic SAMM station system, from dome to fully equipped OTA system. A sentinel may then incorporate (i) magneto-optical filter (MOF) technology and (ii) a coronagraph equipped with a multi-channel spectrograph working in the visible wavelength domain. The technological realisation is associated with the development of the necessary fully automated flare/CME forecast data-processing algorithms. The minimum baseline of a functioning network is a set of SAMM prototypes that consist of a four-strong sentinel station network complemented with the associated fully automated network data processing and control algorithms. **Ideally, these four stations would be separated by about 90 degrees in longitude what would mean a 6-hr long time span of active observation per station. Any longer active observation would result in overlap with the added benefit of cross-checking instrument performances.**




Fig. 15. Potential host sites for future SAMNet sentinels (see Table 2 for further facility details). The blue marks highlight sentinels with about 5-8 hrs time zone differences to indicate where a coronagraph could be added as 4th OTA. Yellow marks are potential future expansion sites.

6.2. The SAMNet sentinel

Each SAMNet station will host a sentinel, a fully-automated telescope that will monitor the Sun using multiple instruments. These sentinels will be connected to form a global network to provide 24-hour solar monitoring. Building on the existing scaled-down SAMM prototype telescope (see Sec. 5.1), here we present an initial design outline for the SAMNet sentinel. Developing a common design simplifies the deployment of new stations in the network, as well as ensuring that all observations will be compatible regardless of which station they originated from.

6.2.1. Sentinel hardware

In order to achieve simultaneous observations of solar active regions at multiple depths, each sentinel will comprise of a single robotic mount with multiple "unit telescopes" (UTs). For sentinels, the components include the various *identical* elements that support the assembly and electro-mechanical stability, powering of the astronomical equipment, supply of resources and the basic conditions for the operation and protection of the observatory (power, tower, dome, weather monitoring, mount, tracking, optics, MOF detectors, camera, data acquisition, processing, data transmission, remote control and automation).

As described in Sec. 5.1, the existing prototype has two unit telescopes with two SAMM MOF units observing the K I and Na D lines. The sentinels used by SAMNet will expand on this to hold at least three such unit telescopes, including the existing K and Na units alongside the proposed Ca MOF under development (see Sec. 5.2.1). The unit telescopes will be aligned to point at the same position, and small guide camera will also be mounted on-axis to ensure the mount reliably tracks the Sun. The potential exists to expand this modular design to include a forth unit telescope, either holding another MOF unit (e.g. He 1083) or a coronagraph-mounted spectrograph.

It is proposed that the SAMNet sentinels will use a similar hardware to their prototype: an array of Celestron OTAs with Baader ND filters, attached to an Astelco mount. The mount may vary subject to the local circumstances at a given station location. By using off-the-shelf hardware as much as possible costs can be reduced and focused on improving the performance of the MOF instruments. The complete SAMNet sentinel node will also include a robotic dome and weather-monitoring stations, to ensure the dome remains closed when the weather is poor.

6.2.2. Control software

Each SAMNet sentinel will be fully-automated, using custom software that autonomously manages the telescope hardware and daily operations. The proposed system will be based on the control system developed for the Gravitational-wave Optical Transient Observer (GOTO) observatory (Dyer et al., 2018). The GOTO project uses very similar hardware design to that proposed for SAMNet: each GOTO mount has eight unit telescopes, and multiple mounts across the world are connected to form a global network. The control software will comprise of multiple independent control daemons which are supervised by a master control program known as the "pilot". An initial design for the core software architecture is shown in Fig. 16.

The primary elements of the control system will be the hardware daemons. A daemon is computer program that runs as a background process, and each category of hardware has a dedicated control daemon. For example, the dome and mount daemons send commands to the dome and

SAMNet team: Solar Activity Monitor Network




Fig. 16. Proposed control software architecture for a SAMNet sentinel.

mount respectively, while the camera daemon centralises commands to all cameras across the three unit telescopes to ensure exposures are synchronised. Each daemon will monitor its associated hardware and then execute commands when received via a command line script, or, in robotic mode, from the pilot. The pilot is the program in overall control of each sentinel when it is operating in robotic mode. During the day the pilot will ensure observations are carried out as required, by issuing commands the hardware daemons. The final program will be the conditions monitor, which continuously monitors local weather stations and other system conditions. Should the weather turn bad or other problems occur the conditions monitor will instruct the pilot to pause and close the dome until it is clear to resume observations.




Fig. 17. A schematic design of multiple SAMNet nodes, expanding on Fig. 16.

6.3. Forming a global network

The central concept of the SAMNet network is to deploy multiple identical sentinels at sites across the Globe. This will make it possible to continuously monitor the Sun for any possible activity, and to issue alerts with as much notice as possible. Each SAMNet site will host one of the sentinels described in Sect. 6.2, and therefore each sentinel is designed to act as a single node in the wider network. As described in Sect. 6.1 least four sites spread equally across the Globe are planned for 24-hour coverage of the Sun, however, any additional sites will provide redundancy for poor weather or other problems. Fig. 15 shows potential host sites proposed by the existing SAMNet partners, each of these sites are listed in Table 2.

Figure 17 shows how each sentinel will transfer images back to the central server for processing. Once the images are transferred, they will be made science-ready and stored at the Sheffield Solar Catalogue (SSC). There, they will be contoured and automatically evaluated by the WG_M method for flare warning. Warning will be issued with probabilities at 6, 12, 18 and 24 hours lead times for a given geographic longitude. The method itself, when applied to photospheric data is capable of forecasting with an about 2-hr lead time at very high accuracy (subject to selection criteria). Using the 3D magnetic skeleton data thanks to a SAMM sentinel output at a given longitude, there would be an additional 2-8 hrs gain in lead time. However, since there is a deployment of SAMMs around the Globe, the lead time may be further improved at a fixed geographic location by a further up to 16 hrs. The total aggregated gain in flare warning would therefore be between 20-26 hrs.

7. Conclusions

The Sun is the main driver of adverse space weather that causes substantial socioeconomic impact at Earth. This includes both direct effects on specific industry sectors (such as electric power, spacecraft and aviation industries) and indirect effects on dependent infrastructures (such as positioning and navigation systems, oil/gas pipelines and financial services). The Lloyd's insurance threat-assessment 2013 report concluded that extreme space weather events would cause USD2.6 trillion damage, and the Space Studies Board, a unit of the National Research Council USA, that full recovery would take 4-10 years (see [here](#)). Superstorms aside, however, the 'nominal' space weather causes an economic loss of Eur10 billion per annum ([Schrijver, 2015](#)).

The network of full 3(+1)-channel SAMM instruments, with SAMNet's cutting-edge MOF and possible coronagraph instrumentation technology, would provide significant improvements to the forecasting of the valiant solar eruptions – the main driver of the extreme space weather events on Earth. This forecast would enable society to take preventive steps to significantly lower the impact of solar eruptions on our technological infrastructure. The mitigation may take place by means based of measuring the LoS components of the solar magnetic and velocity fields in the lower solar atmosphere. In his paper, on page 21, [Beckers \(1971\)](#) says "The magnetic field is however one of the, if not *the*, most important physical quantity in the observable solar atmosphere. Its measurement by any available method is therefore of the greatest importance". SAMNets with its full-scale SAMM sentinel suit is proposed exactly in this spirit. A single SAMM instrument has a relatively low cost and it requires a modest size telescope. Potentially, this could allow a much broader participation in the SAMMNet especially from the Nations with modest budgets for their national science research including countries in African and South American continents. The

instruments could also play a significant role in training the next generation of scientists and the engineers.

SAMNet at its full capacity will yield, for the first time, the foundations of an integrated global space-weather prediction service to mitigate the destructive consequences of space weather. Further, MOF cell spectroscopy has additional wide range of applications from atomic physics, frequency stabilisation in areas such as laser cooling and trapping, high-precision frequency meteorology, remote temperature sensing, atmospheric LIDAR, Doppler velocimetry, optical communications and quantum key distribution in free space, quantum optics, missile defense to navy, to name a few. Most industrial experiments are carried out using the heavier alkali metals, however, there is now a clear rise in applying lighter elements, like Ca.

In the context of other current and future solar observations, SAMM sentinels with their MOF-based sensors and additional coronagraphy instrumentation will also complement data from other space- and ground-based instruments (e.g. the L5 Lagrange mission). In particular, for space-based operational facilities SAMNet will be an ideal, low-cost easy to maintain facility with no telemetry issues.

Obtaining information about lower solar atmospheric magnetic fields, the likely hotbeds of solar eruptions, is often a bottleneck. In particular, there is a clear canopy effect in chromospheric magnetograms, which could be difficult to interpret for modelling input having just LoS observations. With SAMM one will be able to investigate and evaluate how the uncertainties the canopy introduces may effect the accuracy of the forecasting capability.

Acknowledgement RE is grateful to the Science and Technology Facilities Council (STFC, grant numbers ST/M000826/1 and ST/V005979/1) UK and the Royal Society for enabling this research. RE also acknowledges the CAS Presidents International Fellowship Initiative, Grant No. 2019VMA052 and the TOP CLLD 7.1.1-16-H-ERFA-2019-00207-Magyar Napfizikai Alapítvány ID Nr 1-8-2019/P support.

MBK acknowledges STFC grant ST/S000518/1 to Aberystwyth University. MBK also acknowledges the open research program of CAS Key Laboratory of Solar Activity, National Astronomical Observatories, No. KLSA201610. RE and MBK both acknowledge the CAS Key Laboratory of Solar Activity, National Astronomical Observatories Commission for Collaborating Research Program for support received to carry out part of this work. IGH and DP acknowledge support from EPSRC (grant EP/R002061/1). RB acknowledges the support by the Croatian Science Foundation under the project 7549 "Millimeter and submillimeter observations of the solar chromosphere with ALMA". RB, AH, and AMV acknowledge the support from the Austrian-Croatian Bilateral Scientific Project "Comparison of ALMA observations with MHD-simulations of coronal waves interacting with coronal holes". AK and PG acknowledge the support from grant VEGA 2/0048/20. AAP acknowledges NASA 80NSSC18K1242 grant and grant No. 19-02-00088 by RFBR. Sunspot Solar Observatory is a partnership between NSF (1945705), the National Solar Observatory, the state of New Mexico, and consortium partners. TZ acknowledges funding from the Austrian Science Fund (FWF, project P30695-N27). FZ acknowledges support by the Università degli Studi di Catania (Piano per la Ricerca Università di Catania 2016-2018 – Linea di in-

tervento 1 “Chance”; Linea di intervento 2 “Dotazione ordinaria”; Fondi di Ateneo 2020-2022, Università di Catania, Linea Open Access), by the Italian MIUR-PRIN grant 2017APKP7T on “*Circumterrestrial Environment: Impact of Sun-Earth Interaction*”. **YH acknowledges funding from NSFC (11790300,11790301). DK has received funding from the Sêr Cymru II scheme, part-funded by the European Regional Development Fund through the Welsh Government.**

References

- Adams, C. S., and I. G. Hughes. Optics f2f - From Fourier to Fresnel. Oxford University Press, Oxford, UK, 2019. [5.2.1](#)
- Agnelli, G., A. Cacciani, and M. Fofi. The Magneto-Optical Filter. I: Preliminary Observations in Na D Lines. *Sol. Phys.*, **44**(2), 509–518, 1975. [10.1007/BF00153229](#). [5.1.1](#)
- Agnelli, G., A. Cacciani, and M. Fofi. Errata: “The magneto-optical filter. I: Preliminary observations in Na D lines” [Sol. Phys., Vol. 44, p. 509 - 518 (1975)]. *Sol. Phys.*, **46**, 272, 1976. [10.1007/BF00157574](#). [5.1.1](#)
- Anastasiadis, A., A. Papaioannou, I. Sandberg, M. Georgoulis, K. Tziotziou, A. Kouloumvakos, and P. Jiggins. Predicting Flares and Solar Energetic Particle Events: The FORSPEF Tool. *Sol. Phys.*, **292**(9), 134, 2017. [10.1007/s11207-017-1163-7](#). [2](#)
- Avrett, E. H., J. M. Fontenla, and R. Loeser. Formation of the Solar 10830 Angstrom Line. In D. M. Rabin, J. T. Jefferies, and C. Lindsey, eds., *Infrared Solar Physics*, vol. 154, 35, 1994. [3.2](#)
- Baranyi, T., L. Györi, and A. Ludmány. On-line Tools for Solar Data Compiled at the Debrecen Observatory and Their Extensions with the Greenwich Sunspot Data. *Sol. Phys.*, **291**, 3081–3102, 2016. [10.1007/s11207-016-0930-1](#), [1606.00669](#). [2.1](#)
- Barnes, G., K. D. Leka, C. J. Schrijver, T. Colak, R. Qahwaji, et al. A Comparison of Flare Forecasting Methods. I. Results from the All-Clear Workshop. *Astrophys. J.*, **829**, 89, 2016. [10.3847/0004-637X/829/2/89](#), [1608.06319](#). [2](#)
- Beckers, J. M. Narrow band filters based on magneto-optical effects. *Appl. Opt.*, **9**(3), 595, 1970. [10.1364/AO.9.000595](#). [5.1.1](#)
- Beckers, J. M. The Measurement of Solar Magnetic Fields. In R. Howard, ed., *Solar Magnetic Fields*, vol. 43 of *IAU Symposium*, 3, 1971. [7](#)
- Beckers, J. M., and W. J. Wagner. A photographic polarimeter for solar observations. *Appl. Opt.*, **9**, 1933–1934, 1970. [10.1364/AO.9.001933](#). [5.1.1](#)
- Bloom, S. H., P. A. Searcy, K. Choi, R. Kremer, and E. Korevaar. Helicopter plume detection by using an ultranarrow-band noncoherent laser Doppler velocimeter. *Opt. Lett.*, **18**(3), 244–246, 1993. [10.1364/OL.18.000244](#). [3.1](#)
- Bobra, M. G., and S. Couvidat. Solar Flare Prediction Using SDO/HMI Vector Magnetic Field Data with a Machine-learning Algorithm. *Astrophys. J.*, **798**(2), 135, 2015. [10.1088/0004-637X/798/2/135](#), [1411.1405](#). [2](#)

- 829 Cacciani, A., M. Cimino, and M. Fofi. A Short Report on the Magnetic Beam Absorption Filter Research
830 at the Rome Astronomical Observatory. In R. Howard, ed., *Solar Magnetic Fields*, vol. 43 of *IAU*
831 *Symposium*, 94, 1971. 5.1.1
- 832 Cacciani, A., and M. Fofi. The Magneto-Optical Filter. II. Velocity Field Measurements. *Sol. Phys.*, **59**(1),
833 179–189, 1978. 10.1007/BF00154941. 5.1.1
- 834 Cacciani, A., C. Marmolino, P. F. Moretti, M. Oliviero, G. Severino, and L. A. Smaldone. Simultaneous
835 Doppler and magnetic solar maps from a MOF installed at the Osservatorio di Capodimonte.
836 *Mem. Soc. Astron. Italiana*, **68**, 467, 1997a. 3.1, 3.2, 5.1.1
- 837 Cacciani, A., and P.-F. Moretti. Magneto-optical filter: concept and applications in astronomy. In D. L.
838 Crawford and E. R. Craine, eds., *Proc. SPIE*, vol. 2198 of *Society of Photo-Optical Instrumentation*
839 *Engineers (SPIE) Conference Series*, 219–228, 1994. 10.1117/12.176751. 3.1
- 840 Cacciani, A., P. F. Moretti, and W. E. Rodgers. Measuring Doppler and Magnetic Fields Simultaneously.
841 *Sol. Phys.*, **174**, 115–128, 1997b. 10.1023/A:1004935524038. 3.1, 3.2
- 842 Cacciani, A., D. Ricci, P. Rosati, E. J. Rhodes, E. Smith, S. Tomczyk, and R. K. Ulrich. Solar magnetic fields
843 measurements with a magneto-optical filter. *Nuovo Cimento C Geophysics Space Physics C*, **13**, 125–130,
844 1990. 10.1007/BF02515781. 3.2
- 845 Campi, C., F. Benvenuto, A. M. Massone, D. S. Bloomfield, M. K. Georgoulis, and M. Piana. Feature
846 Ranking of Active Region Source Properties in Solar Flare Forecasting and the Uncompromised
847 Stochasticity of Flare Occurrence. *Astrophys. J.*, **883**(2), 150, 2019. [http://dx.doi.org/10.3847/](http://dx.doi.org/10.3847/1538-4357/ab3c26)
848 [1538-4357/ab3c26](http://dx.doi.org/10.3847/1538-4357/ab3c26), 1906.12094. 2
- 849 Camporeale, E. The Challenge of Machine Learning in Space Weather: Nowcasting and Forecasting. *Space*
850 *Weather*, **17**(8), 1166–1207, 2019. 10.1029/2018SW002061, URL [https://agupubs.onlinelibrary.](https://agupubs.onlinelibrary.wiley.com/doi/abs/10.1029/2018SW002061)
851 [wiley.com/doi/abs/10.1029/2018SW002061](https://agupubs.onlinelibrary.wiley.com/doi/abs/10.1029/2018SW002061). 2, 2.2
- 852 Chang, P., Y. Chen, H. Shang, X. Guan, H. Guo, J. Chen, and B. Luo. A Faraday laser operating on Cs 852 nm
853 transition. *APPLIED PHYSICS B-LASERS AND OPTICS*, **125**(12), 2019. 10.1007/s00340-019-7342-5.
854 3.1
- 855 Cimino, M., A. Cacciani, and M. Fofi. Some Developments of the Magnetic Beam Absorption Filter.
856 *Sol. Phys.*, **11**(2), 319–333, 1970. 10.1007/BF00155231. 3.2, 5.1.1
- 857 Cimino, M., A. Cacciani, and N. Sopranzi. An Instrument to measure Solar Magnetic Fields by an Atomic-
858 Beam Method. *Sol. Phys.*, **3**(4), 618–622, 1968. 10.1007/BF00151943. 3.1, 3.2, 5.1.1
- 859 Cui, Y., R. Li, L. Zhang, Y. He, and H. Wang. Correlation Between Solar Flare Productivity and Photospheric
860 Magnetic Field Properties. 1. Maximum Horizontal Gradient, Length of Neutral Line, Number of Singular
861 Points. *Sol. Phys.*, **237**, 45–59, 2006. 10.1007/s11207-006-0077-6. 2.1
- 862 DeForest, C. E., H. J. Hagenaar, D. A. Lamb, C. E. Parnell, and B. T. Welsch. Solar Magnetic Tracking. I.
863 Software Comparison and Recommended Practices. *Astrophys. J.*, **666**, 576–587, 2007. 10.1086/518994,
864 0704.2921. 2.3.1
- 865 Deng, Y. Y. Introduction to the Chinese Giant Solar Telescope. In *Astronomical Society of India Conference*
866 *Series*, vol. 2 of *Astronomical Society of India Conference Series*, 31–36, 2011. 6.1

- 867 Dyer, M. J., V. S. Dhillon, S. Littlefair, D. Steeghs, K. Ulaczyk, P. Chote, D. Galloway, and E. Rol. A
868 telescope control and scheduling system for the Gravitational-wave Optical Transient Observer (GOTO).
869 In Proceedings of the SPIE, vol. 10704 of Society of Photo-Optical Instrumentation Engineers (SPIE)
870 Conference Series, 107040C, 2018. 10.1117/12.2311865, [1807.01614](#). [6.2.2](#)
- 871 Eastwood, J. P., E. Biffis, M. A. Hapgood, L. Green, M. M. Bisi, R. D. Bentley, R. Wicks, L. A. McKinnell,
872 M. Gibbs, and C. Burnett. The Economic Impact of Space Weather: Where Do We Stand? Risk Analysis,
873 **37**(2), 206–218, 2017. 10.1111/risa.12765. [1](#)
- 874 Finsterle, W., S. M. Jefferies, A. Cacciani, P. Rapex, and S. W. McIntosh. Helioseismic Mapping of
875 the Magnetic Canopy in the Solar Chromosphere. Astrophys. J. Lett., **613**(2), L185–L188, 2004a.
876 10.1086/424996. [5.1.1](#)
- 877 Finsterle, W., S. M. Jefferies, A. r. Cacciani, P. Rapex, C. Giebink, A. Knox, and V. Dimartino. Seismology
878 of the solar atmosphere. Sol. Phys., **220**(2), 317–331, 2004b. 10.1023/B:SOLA.0000031397.73790.7b.
879 [3.2](#), [5.1.1](#)
- 880 Florios, K., I. Kontogiannis, S.-H. Park, J. A. Guerra, F. Benvenuto, D. S. Bloomfield, and M. K. Georgoulis.
881 Forecasting Solar Flares Using Magnetogram-based Predictors and Machine Learning. Sol. Phys., **293**(2),
882 28, 2018. 10.1007/s11207-018-1250-4, [1801.05744](#). [2](#)
- 883 Forte, R., S. Jefferies, E. Pietropaolo, S. Scardigli, L. Giovannelli, D. Del Moro, and F. Berrilli. MOTH II
884 calibration pipeline and data merging with SDO/HMI and SDO/AIA. In SOLARNET IV: The Physics of
885 the Sun from the Interior to the Outer Atmosphere, 122, 2017. [3.2](#)
- 886 Forte, R., S. M. Jefferies, F. Berrilli, D. Del Moro, B. Fleck, L. Giovannelli, N. Murphy, E. Pietropaolo,
887 and W. Rodgers. The MOTH II Doppler-Magnetographs and Data Calibration Pipeline. In C. Foullon
888 and O. E. Malandraki, eds., Space Weather of the Heliosphere: Processes and Forecasts, vol. 335 of IAU
889 Symposium, 335–339, 2018. 10.1017/S1743921318000029. [3.2](#)
- 890 Gary, G. A. Linear force-free magnetic fields for solar extrapolation and interpretation.
891 Astrophys. J. Suppl. Ser., **69**, 323–348, 1989. 10.1086/191316. [2.3.1](#)
- 892 Georgoulis, M. K. Toward an Efficient Prediction of Solar Flares: Which Parameters, and How? Entropy,
893 **15**(11), 5022–5052, 2013. 10.3390/e15115022. [2](#)
- 894 Guevara Gómez, J. C., J. C. Martínez Oliveros, and B. Calvo-Mozo. First Colombian Solar Radio
895 Interferometer: current stage. In S. Vargas Domínguez, A. G. Kosovichev, P. Antolin, and L. Harra,
896 eds., Fine Structure and Dynamics of the Solar Atmosphere, vol. 327 of IAU Symposium, 16–19, 2017.
897 10.1017/S1743921317000084. [2](#)
- 898 Haberreiter, M., W. Finsterle, and S. M. Jefferies. On the observation of traveling acoustic waves in
899 the solar atmosphere using a magneto-optical filter. Astronomische Nachrichten, **328**(3), 211, 2007.
900 10.1002/asna.200610721. [5.1.1](#)
- 901 Hasan, S. S., D. Soltau, H. Kärcher, M. Süß, and T. Berkefeld. NLST: India’s National Large Solar Telescope.
902 Astronomische Nachrichten, **331**(6), 628, 2010. 10.1002/asna.201011389. [6.1](#)
- 903 Huang, W., X. Chu, J. Wiig, B. Tan, C. Yamashita, et al. Field demonstration of simultaneous wind and
904 temperature measurements from 5 to 50 km with a Na double-edge magneto-optic filter in a multi-frequency
905 Doppler lidar. Opt. Lett., **34**(10), 1552–1554, 2009. 10.1364/OL.34.001552. [3.1](#)

- Huang, X., H. Wang, L. Xu, J. Liu, R. Li, and X. Dai. Deep Learning Based Solar Flare Forecasting Model. I. Results for Line-of-sight Magnetograms. *Astrophys. J.*, **856**(1), 7, 2018. 10.3847/1538-4357/aae00. 2.2
- Keaveney, J., W. J. Hamlyn, C. S. Adams, and I. G. Hughes. A single-mode external cavity diode laser using an intra-cavity atomic Faraday filter with short-term linewidth <400 kHz and long-term stability of <1 MHz. *Review of Scientific Instruments*, **87**(9), 095,111, 2016. 10.1063/1.4963230. 3.1
- Keaveney, J., S. A. Wrathmall, C. S. Adams, and I. G. Hughes. Optimized ultra-narrow atomic bandpass filters via magneto-optic rotation in an unconstrained geometry. *Optics Letters*, **43**(17), 4272, 2018. 10.1364/OL.43.004272, 1806.08705. 5.2.1
- Kim, T., E. Park, H. Lee, Y.-J. Moon, S.-H. Bae, et al. Solar farside magnetograms from deep learning analysis of STEREO/EUVI data. *Nature Astronomy*, **3**, 397–400, 2019. 10.1038/s41550-019-0711-5. 2
- Korsós, M. B., P. Chatterjee, and R. Erdélyi. Applying the Weighted Horizontal Magnetic Gradient Method to a Simulated Flaring Active Region. *Astrophys. J.*, **857**, 103, 2018. 10.3847/1538-4357/aab891, 1804.10351. 2.3.2
- Korsós, M. B., M. K. Georgoulis, N. Gyenge, S. K. Bisoi, S. Yu, S. Poedts, C. J. Nelson, J. Liu, Y. Yan, and R. Erdélyi. Solar Flare Prediction Using Magnetic Field Diagnostics above the Photosphere. *Astrophys. J.*, **896**(2), 119, 2020a. 10.3847/1538-4357/ab8fa2, 2005.12180. 2.1, 2.3, 2.3.1, 2.3.2, 2.3.3
- Korsós, M. B., A. Ludmány, R. Erdélyi, and T. Baranyi. On Flare Predictability Based on Sunspot Group Evolution. *Astrophys. J. Lett.*, **802**(2), L21, 2015. 10.1088/2041-8205/802/2/L21, 1503.04634. 2.1, 2.1
- Korsós, M. B., P. Romano, H. Morgan, Y. Ye, R. Erdélyi, and F. Zuccarello. Differences in Periodic Magnetic Helicity Injection Behavior between Flaring and Non-flaring Active Regions: Case Study. *Astrophys. J. Lett.*, **897**(2), L23, 2020b. 10.3847/2041-8213/ab9d7a, 2006.07659. 1
- Korsós, M. B., S. Yang, and R. Erdélyi. Investigation of pre-flare dynamics using the weighted horizontal magnetic gradient method: From small to major flare classes. *Journal of Space Weather and Space Climate*, **9**, A6, 2019. 10.1051/swsc/2019002, 1901.05984. 2.1, 2.1, 2.3, 2.3.2
- Kuridze, D., M. Mathioudakis, D. J. Christian, A. F. Kowalski, D. B. Jess, S. D. T. Grant, T. Kawate, P. J. A. Simões, J. C. Allred, and F. P. Keenan. Observations and Simulations of the Na I D₁ Line Profiles in an M-class Solar Flare. *Astrophys. J.*, **832**(2), 147, 2016. 10.3847/0004-637X/832/2/147, 1609.08120. 3.2, 3.2
- Leenaarts, J., T. Golding, M. Carlsson, T. Libbrecht, and J. Joshi. The cause of spatial structure in solar He I 1083 nm multiplet images. *Astron. Astrophys.*, **594**, A104, 2016. 10.1051/0004-6361/201628490, 1608.00838. 3.2
- Leka, K. D., G. Barnes, and E. Wagner. The NWRA Classification Infrastructure: description and extension to the Discriminant Analysis Flare Forecasting System (DAFFS). *J. Space Weather and Space Clim.*, **8**, A25, 2018. <http://dx.doi.org/10.1051/swsc/2018004>, 1802.06864. 2
- Leka, K. D., S.-H. Park, K. Kusano, J. Andries, G. Barnes, et al. A Comparison of Flare Forecasting Methods. II. Benchmarks, Metrics, and Performance Results for Operational Solar Flare Forecasting Systems. *Astrophys. J. Suppl. Ser.*, **243**(2), 36, 2019a. 10.3847/1538-4365/ab2e12, 1907.02905. 2

SAMNet team: Solar Activity Monitor Network

- 944 Leka, K. D., S.-H. Park, K. Kusano, J. Andries, G. Barnes, et al. A Comparison of Flare Forecasting Methods.
945 III. Systematic Behaviors of Operational Solar Flare Forecasting Systems. *Astrophys. J.*, **881**(2), 101,
946 2019b. 10.3847/1538-4357/ab2e11, [1907.02909](#). [2](#)
- 947 Lide, D. R. CRC handbook of chemistry and physics: a ready-reference book of chemical and physical data.
948 CRC press, 1995. [5.2.1](#)
- 949 Liu, J., Y. Wang, X. Huang, M. B. Korsós, Y. Jiang, Y. Wang, and R. Erdélyi. Reliability of AI-generated
950 magnetograms from only EUV images. *Nature Astronomy*, **5**, 108–110, 2021. 10.1038/s41550-021-
951 01310-6. [2](#)
- 952 Liu, J., Y. Ye, C. Shen, Y. Wang, and R. Erdélyi. A New Tool for CME Arrival Time Prediction using Machine
953 Learning Algorithms: CAT-PUMA. *Astrophys. J.*, **855**(2), 109, 2018. 10.3847/1538-4357/aaae69, [1802.](#)
954 [02803](#). [6.1](#)
- 955 Liu, Z., J. Xu, B.-Z. Gu, S. Wang, J.-Q. You, et al. New vacuum solar telescope and observations with high
956 resolution. *Research in Astronomy and Astrophysics*, **14**(6), 705-718, 2014. 10.1088/1674-4527/14/6/009.
957 [6.1](#)
- 958 Miao, X., L. Yin, W. Zhuang, B. Luo, A. Dang, J. Chen, and H. Guo. Note: Demonstration of an
959 external-cavity diode laser system immune to current and temperature fluctuations. *Review of Scientific*
960 *Instruments*, **82**(8), 086,106, 2011. 10.1063/1.3624696, <https://doi.org/10.1063/1.3624696>. [3.1](#)
- 961 Moretti, P. F., F. Berrilli, A. Bigazzi, S. M. Jefferies, N. Murphy, L. Roselli, and M. P. di Mauro. Future
962 instrumentation for solar physics: a double channel MOF imager on board ASI Space Mission ADAHELI.
963 *Astrophys. Space Sci.*, **328**(1-2), 313–318, 2010. 10.1007/s10509-009-0251-z. [3.2](#)
- 964 Moretti, P. F., G. Severino, G. Cauzzi, K. Reardon, T. Straus, A. Cacciani, C. Marmolino, M. Oliviero, and
965 L. A. Smaldone. The Magneto-Optical Filter in Napoli: Perspectives and Test Observations, vol. 225 of
966 *Astrophysics and Space Science Library*, 293–296, 1997. 10.1007/978-94-011-5167-2-32. [5.1.1](#)
- 967 Murphy, N., E. Smith, W. Rodgers, and S. Jefferies. Chromospheric Observations in the Helium 1083nm
968 Line - A new Instrument. In B. Fleck, T. H. Zurbuchen, and H. Lacoste, eds., *Solar Wind 11/SOHO 16,*
969 *Connecting Sun and Heliosphere*, vol. 592 of *ESA Special Publication*, 511, 2005. [3.2](#), [5.2](#)
- 970 Nishizuka, N., K. Sugiura, Y. Kubo, M. Den, and M. Ishii. Deep Flare Net (DeFN) Model for Solar Flare
971 Prediction. *Astrophys. J.*, **858**(2), 113, 2018. 10.3847/1538-4357/aab9a7, [1805.03421](#). [2.2](#)
- 972 Oliviero, M., M. Dolci, G. Severino, T. Straus, A. Cacciani, and P. F. Moretti. VAMOS: Velocity and Magnetic
973 Observations of the Sun. In F.-L. Deubner, J. Christensen-Dalsgaard, and D. Kurtz, eds., *New Eyes to See*
974 *Inside the Sun and Stars*, vol. 185 of *IAU Symposium*, 53, 1998a. [5.1.1](#)
- 975 Oliviero, M., P. F. Moretti, G. Severino, T. Straus, M. Magrì, and A. Tripicchio. Preliminary Results
976 on the Solar Photospheric Dynamics Observed with Vamos. *Sol. Phys.*, **209**(1), 21–35, 2002.
977 10.1023/A:1020904204842. [3.2](#), [5.1.1](#), [12](#), [5.1.1](#)
- 978 Oliviero, M., G. Severino, and T. Straus. The VAMOS Data Analysis Pipeline. In S. Korzennik, ed., *Structure*
979 *and Dynamics of the Interior of the Sun and Sun-like Stars*, vol. 418 of *ESA Special Publication*, 275,
980 1998b. [5.1.1](#)

- 981 Park, E., Y.-J. Moon, S. Shin, K. Yi, D. Lim, H. Lee, and G. Shin. Application of the Deep Convolutional
982 Neural Network to the Forecast of Solar Flare Occurrence Using Full-disk Solar Magnetograms.
983 Astrophys. J., **869**(2), 91, 2018. 10.3847/1538-4357/aaed40. [2.2](#)
- 984 Pevtsov, A. A. Space Weather Forecasting and Supporting Research in the USA. Geomagnetism and
985 Aeronomy, **57**(7), 769–775, 2017. 10.1134/S0016793217070179. [1](#)
- 986 Pötzi, W., A. M. Veronig, G. Riegler, U. Amerstorfer, T. Pock, M. Temmer, W. Polanec, and D. J.
987 Baumgartner. Real-time Flare Detection in Ground-Based H α Imaging at Kanzelhöhe Observatory.
988 Sol. Phys., **290**(3), 951–977, 2015. 10.1007/s11207-014-0640-5, [1411.3896](#). [2](#)
- 989 Pötzi, W., A. M. Veronig, and M. Temmer. An Event-Based Verification Scheme for the Real-Time Flare
990 Detection System at Kanzelhöhe Observatory. Sol. Phys., **293**(6), 94, 2018. 10.1007/s11207-018-1312-7.
991 [2](#)
- 992 Pötzi, W., A. M. Veronig, M. Temmer, D. J. Baumgartner, H. Freislich, and H. Strutzmann. 70 Years
993 of Sunspot Observations at the Kanzelhöhe Observatory: Systematic Study of Parameters Affecting the
994 Derivation of the Relative Sunspot Number. Sol. Phys., **291**(9-10), 3103–3122, 2016. 10.1007/s11207-
995 016-0857-6, [1512.00270](#). [2](#)
- 996 Qahwaji, R., and T. Colak. Automatic Short-Term Solar Flare Prediction Using Machine Learning and
997 Sunspot Associations. Sol. Phys., **241**(1), 195–211, 2007. 10.1007/s11207-006-0272-5. [2](#)
- 998 Quintero Noda, C., H. Uitenbroek, Y. Katsukawa, T. Shimizu, T. Oba, et al. Solar polarimetry through
999 the K I lines at 770 nm. Mon. Not. R. Astron. Soc., **470**(2), 1453–1461, 2017. 10.1093/mnras/stx1344,
1000 [1705.10002](#). [3.2](#)
- 1001 Ravindra, B., P. Kesavan, K. C. Thulasidharen, M. Rajalingam, K. Sagayanathan, et al. Installation of solar
1002 chromospheric telescope at the Indian Astronomical Observatory, Merak. Journal of Astrophysics and
1003 Astronomy, **39**(5), 60, 2018. 10.1007/s12036-018-9554-1. [6.1](#)
- 1004 Rhodes, J., E. J., A. Cacciani, S. Tomczyk, R. K. Ulrich, J. Blamont, R. F. Howard, P. Dumont, and E. J.
1005 Smith. A compact dopplergraph/magnetograph suitable for space-based measurements of solar oscillations
1006 and magnetic fields. Advances in Space Research, **4**(8), 103–112, 1984. 10.1016/0273-1177(84)90371-5.
1007 [5.1.1](#)
- 1008 Schrijver, C. J. A Characteristic Magnetic Field Pattern Associated with All Major Solar Flares and Its Use
1009 in Flare Forecasting. Astrophys. J. Lett., **655**, L117–L120, 2007. 10.1086/511857. [2.1](#)
- 1010 Schrijver, C. J. Socio-Economic Hazards and Impacts of Space Weather: The Important Range Between
1011 Mild and Extreme. Space Weather, **13**(9), 524–528, 2015. 10.1002/2015SW001252, [1507.08730](#). [7](#)
- 1012 Severino, G., P. F. Moretti, M. Oliviero, and Vamos Team. The Velocity And Magnetic Observations of the
1013 Sun (VAMOS) project: status and future prospects. In A. Wilson and P. L. Pallé, eds., SOHO 10/GONG
1014 2000 Workshop: Helio- and Asteroseismology at the Dawn of the Millennium, vol. 464 of ESA Special
1015 Publication, 337–340, 2001. [5.1.1](#)
- 1016 Shan, X., X. Sun, J. Luo, Z. Tan, and M. Zhan. Free-space quantum key distribution with Rb vapor filters.
1017 Applied Physics Letters, **89**(19), 191,121, 2006. 10.1063/1.2387867, [https://doi.org/10.1063/1.](https://doi.org/10.1063/1.2387867)
1018 [2387867](#). [3.1](#)

- Stangalini, M., P. F. Moretti, F. Berrilli, D. Del Moro, S. M. Jefferies, G. Severino, and M. Oliviero. DIMMI-2h a MOF-based instrument for Solar Satellite ADAHELI. In Proc. SPIE, vol. 8148 of Society of Photo-Optical Instrumentation Engineers (SPIE) Conference Series, 81480U, 2011. 10.1117/12.893579. [5.1.1](#)
- Stangalini, M., R. Piazzesi, R. Speziali, and L. Dal Sasso. SAMM: the solar activity MOF monitor. In Proc. SPIE, vol. 10700 of Society of Photo-Optical Instrumentation Engineers (SPIE) Conference Series, 107001K, 2018. 10.1117/12.2313373. [5.1.1](#), [5.1.1](#)
- Supriya, H. D., H. N. Smitha, K. N. Nagendra, J. O. Stenflo, M. Bianda, R. Ramelli, B. Ravindra, and L. S. Anusha. Center-to-limb Observations and Modeling of the Ca I 4227 Å Line. Astrophys. J., **793**(1), 42, 2014. 10.1088/0004-637X/793/1/42, [1407.5461](#). [3.2](#), [5.2.1](#)
- Tomczyk, S., A. Cacciani, and S. A. Veitzer. LOWL - an Instrument to Observe Low-Degree Solar Oscillations. In T. M. Brown, ed., GONG 1992. Seismic Investigation of the Sun and Stars, vol. 42 of Astronomical Society of the Pacific Conference Series, 469, 1993. [5.1.1](#)
- Tomczyk, S., K. Streander, G. Card, D. Elmore, H. Hull, and A. Cacciani. An Instrument to Observe Low-Degree Solar Oscillations. Sol. Phys., **159**(1), 1–21, 1995. 10.1007/BF00733027. [5.1.1](#)
- Uitenbroek, H. Chromospheric Diagnostics. In J. Leibacher, R. F. Stein, and H. Uitenbroek, eds., Solar MHD Theory and Observations: A High Spatial Resolution Perspective, vol. 354 of Astronomical Society of the Pacific Conference Series, 313, 2006. [3.2](#)
- Veronig, A. M., and W. Pötzi. Ground-based Observations of the Solar Sources of Space Weather. In I. Dorotovic, C. E. Fischer, and M. Temmer, eds., Coimbra Solar Physics Meeting: Ground-based Solar Observations in the Space Instrumentation Era, vol. 504 of Astronomical Society of the Pacific Conference Series, 247, 2016. [1602.02721](#). [2](#)
- Wallace, L., K. H. Hinkle, W. C. Livingston, and S. P. Davis. An Optical and Near-infrared (2958–9250 Å) Solar Flux Atlas. Astrophys. J. Suppl. Ser., **195**(1), 6, 2011. 10.1088/0067-0049/195/1/6. [11](#)
- Wang, J., Y. Zhang, S. A. Hess Webber, S. Liu, X. Meng, and T. Wang. Solar Flare Predictive Features Derived from Polarity Inversion Line Masks in Active Regions Using an Unsupervised Machine Learning Algorithm. Astrophys. J., **892**(2), 140, 2020. 10.3847/1538-4357/ab7b6c. [2.2](#)
- Wang, Y., J. Liu, Y. Jiang, and R. Erdélyi. CME Arrival Time Prediction Using Convolutional Neural Network. Astrophys. J., **881**(1), 15, 2019. 10.3847/1538-4357/ab2b3e. [2](#), [6.1](#)
- Welsch, B. T., and D. W. Longcope. Magnetic Helicity Injection by Horizontal Flows in the Quiet Sun. I. Mutual-Helicity Flux. Astrophys. J., **588**, 620–629, 2003. 10.1086/368408. [2.3.1](#)
- Wiegmann, T., and T. Sakurai. Solar Force-free Magnetic Fields. Living Reviews in Solar Physics, **9**, 2012. 10.12942/lrsp-2012-5, [1208.4693](#). [2](#)
- Xia, Y., X. Cheng, F. Li, Y. Yang, X. Lin, J. Jiao, L. Du, J. Wang, and G. Yang. Sodium lidar observation over full diurnal cycles in Beijing, China. Appl. Opt., **59**(6), 1529, 2020. 10.1364/AO.382077. [3.1](#)
- Yan, Y., Z. Chen, W. Wang, F. Liu, L. Geng, L. Chen, C. Tan, X. Chen, C. Su, and B. Tan. Mingantu Spectral Radiograph for Solar and Space Weather Studies. Front. Astron. Space Sci., **8**(584043), 1–13, 2021. 10.3389/fspas.2021.584043, URL <https://www.frontiersin.org/articles/10.3389/fspas.2021.584043/abstract>. [6.1](#)

SAMNet team: Solar Activity Monitor Network

- 1058 Yan, Y., W. Wang, L. Chen, F. Liu, L. Geng, and Z. Chen. New Interplanetary Scintillation Array in China
1059 for Space Weather. Sun and Geosphere, **13**, 153–155, 2018. 10.31401/SunGeo.2018.02.05. [6.1](#)
- 1060 Yan, Y., J. Zhang, W. Wang, F. Liu, Z. Chen, and G. Ji. The Chinese Spectral Radioheliograph—CSRH.
1061 Earth Moon and Planets, **104**(1-4), 97–100, 2009. 10.1007/s11038-008-9254-y. [6.1](#)
- 1062 Yang, Y., X. Cheng, F. Li, X. Hu, X. Lin, and S. Gong. A flat spectral Faraday filter for sodium lidar. Optics
1063 Letters, **36**(7), 1302, 2011. 10.1364/OL.36.001302. [3.1](#), [3.1](#), [4](#)
- 1064 Ye, Y., M. B. Korsós, and R. Erdélyi. Detailed analysis of dynamic evolution of three Active Regions at the
1065 photospheric level before flare and CME occurrence. Advances in Space Research, **61**(2), 673–682, 2018.
1066 10.1016/j.asr.2017.09.038. [1](#)
- 1067 Zhao, J., D. Hing, R. Chen, and S. Hess Webber. Imaging the Sun’s Far-side Active Regions by Applying
1068 Multiple Measurement Schemes on Multiskip Acoustic Waves. Astrophys. J., **887**(2), 216, 2019.
1069 10.3847/1538-4357/ab5951, [1912.06736](#). [1](#)
- 1070 Zielińska, J. A., F. A. Beduini, V. G. Lucivero, and M. W. Mitchell. Atomic filtering for hybrid
1071 continuous-variable/discrete-variable quantum optics. Opt. Express, **22**(21), 25,307–25,317, 2014.
1072 10.1364/OE.22.025307. [3.1](#)